



Anatomy of the lung revisited by 3D-CT imaging

Seshiru Nakazawa^{1,2^}, Toshiteru Nagashima¹, Natsuko Kawatani¹, Patrick C. Gedeon³,
Ariadne K. DeSimone⁴, Hitoshi Igai⁵, Takayuki Kosaka^{1,6}, Ken Shirabe¹

¹Department of General Surgical Science, Gunma University Graduate School of Medicine, Maebashi, Japan; ²Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA, USA; ³Department of Surgery, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA; ⁴Department of Radiology, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA; ⁵Department of General Thoracic Surgery, Japanese Red Cross Maebashi Hospital, Maebashi, Japan; ⁶Department of Thoracic Surgery, National Hospital Organization Takasaki General Medical Center, Takasaki, Japan

Contributions: (I) Conception and design: S Nakazawa, H Igai, PC Gedeon, AK DeSimone, T Kosaka; (II) Administrative support: S Nakazawa, T Nagashima, K Shirabe; (III) Provision of study materials or patients: S Nakazawa, T Nagashima, N Kawatani; (IV) Collection and assembly of data: S Nakazawa, T Nagashima, N Kawatani, T Kosaka; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Seshiru Nakazawa, MD PhD. Department of General Surgical Science, Gunma University Graduate School of Medicine, 3-39-22 Showa-machi, Maebashi, 371-8511, Japan. Email: snakazawa@gunma-u.ac.jp.

Abstract: The anatomy of the lung was originally described based on data acquired from cadaveric studies and surgical findings. Over time, computed tomography (CT) and three-dimensional (3D) imaging techniques have been developed, allowing for reconstruction and understanding of lung anatomy in a more intuitive way. The wide adoption of 3D-CT imaging technology has led to a variety of anatomical studies performed not only by anatomists but also by surgeons and radiologists. Such studies have led to new or modified classification systems, shed light on lung anatomy from a useful surgical viewpoint, and enabled us to analyze lung anatomy with a focus on particular anatomical features. 3D images also allow for enhanced pre- and intra-operative simulation, improved surgical safety, enhanced educational utility, and the capacity to perform large-scale anatomical studies in shorter time frames. We will review here the key features of 3D-CT imaging of the lung, along with representative anatomical studies regarding (I) general lung anatomy, (II) anatomy of the right and left lobes, and (III) features of interlobar vessels. The current surge of 3D imaging analysis shows that the field is growing, with the technology continuing to improve. Future studies using these new and innovative methodologies will continue to refine our understanding of lung anatomy while enhancing our ability to perform safe and effective surgical resections.

Keywords: Lung; anatomy; 3D imaging

Received: 16 February 2023; Accepted: 11 May 2023; Published online: 24 May 2023.

doi: 10.21037/vats-23-21

View this article at: <https://dx.doi.org/10.21037/vats-23-21>

Introduction

The anatomy of the lung has been studied in detail since the 1930's with an international nomenclature proposed in 1950 (1). Knowledge of lung segments has been periodically revised, mostly based on input from cadaveric studies and surgical findings (2-5). This led to a textbook presented

by Yamashita in 1978 which today remains widely used and a key reference in many publications (5). However, information collected from cadavers has drawbacks including postmortem modifications or limitations in analysis due to difficulties procuring cases. Since these initial cadaveric and surgical studies, non-invasive

[^] 0000-0001-6218-9396.

Table 1 Features of 3D-CT imaging in lung

-
- Better recognition of anatomical structures compared to two-dimensional images
 - Quick and intuitive recognition of the anatomy and anomalies
 - Pre- and intra-operative simulation
 - Improved surgical safety
 - Enhanced educational utility
 - New types of analyses involving lung volume or lung parenchyma
 - Large-scale studies with short study times (>1,000 cases)
-

3D, three-dimensional; CT, computed tomography.

imaging technology has flourished enabling analyses from a radiological perspective. In particular, computed tomography (CT) and three-dimensional (3D) imaging has made it possible to reconstruct the anatomy of the lung allowing for a more intuitive understanding of the spatial relationships between structures (6). 3D-CT imaging involves postprocessing of the entire multidetector CT data set to generate 3D volume-rendered images, allowing for visualization and manipulation of objects represented as sample data in 3D. The adoption of 3D-CT imaging has led to a variety of analyses led not only by anatomists but also by surgeons and radiologists. Thoracic surgeons also became actively engaged in anatomical analysis using 3D imaging, partly driven by the surgical need to better understand the anatomy of the lung (7-12). We will review here key features of 3D-CT imaging of the lung and present representative anatomical studies based on 3D-CT images.

Features of 3D-CT Imaging

A major advantage of 3D-CT imaging lies in its ability to allow for better recognition of anatomical structures compared to two-dimensional (2D) images (*Table 1*). Although conventional CT images do include information necessary to analyze anatomy, they are not always sufficient to fully perceive the spatial relationships between anatomical structures. A trained physician could recreate anatomical structures in his or her mind (13), but the technique requires skill and is time consuming. 3D-CT images enable surgeons to quickly and more intuitively recognize the anatomy and associated anomalies (14). Also, images can be shared for educational purposes or used for pre- and intra-operative simulation (10).

Another benefit of 3D-CT imaging is the associated improvement in surgical safety. With the increasing number of segmentectomies being performed, a detailed understanding of each patient's unique anatomy, which includes the spatial relationship between bronchial, vascular, and parenchymal structures, is of increasing importance. In 2011, Oizumi *et al.* presented the usefulness of preoperative 3D-CT imaging in thoracoscopic segmentectomy and reported a 98% success rate (8). Many others have also reported the benefits of 3D-CT imaging in lung surgery (15). Surgeons performing segmentectomy are also more likely to be faced with segmental variations, and some commonly encountered variations have been analyzed in more detail, such as the subsuperior segment (16-18). Accordingly, an understanding of segmental anatomy and associated anatomic variants is essential for segmentectomy (*Figure 1*). However, a basic knowledge of anatomical variations is also beneficial when performing a routine lobectomy. We should always bear in mind the possibility of encountering common variations, such as an anomalous V^2 , a mediastinal lingular artery, or a lingular vein draining into the inferior pulmonary vein, each of which may result in surgical complications should they fail to be appreciated.

The use of 3D-CT imaging furthermore allows for new types of analysis such as the relationship between lung volume and segmental anatomy (19) or the spatial relationship between intersegmental planes and intersegmental veins (20). Mimae *et al.* reported that the main root of the intersegmental vein (V^3a+b) between the upper and lingular divisions was always located in the upper division, whereas the root of the intersegmental vein (V^6b+c) between S^6 and basal segments was always located in the basal segment; it is important to know that intersegmental veins are not always located on the intersegmental plane when dividing the lung parenchyma along these intersegmental veins (20).

Also, several 3D-CT imaging studies have enrolled a large number of patients within a very short amount of time, some including more than 1,000 cases and others occasionally exceeding 5,000 cases (17,21-23). Such large-scale analyses would be difficult in a cadaveric study. Furthermore, classifying all cases into diverse anatomical categories with 2D images alone would also be an immensely complicated task. Accordingly, the volume, speed, and rigor at which complex anatomic studies may be conducted and analyzed using 3D imaging provides distinct advantages over studies using cadaveric specimens or 2D images alone.




Figure 1 Schema and nomenclature of bronchopulmonary segments. The medial basal segment (S⁷) and subsuperior segment (S*) are not depicted in the schema.

General anatomy of the lung

In 2010, Akiba *et al.* analyzed variations of the pulmonary vein using 3D-CT images (Table 2) (24). They reported that most patients had the expected anatomy (98% of the left side and 86% of the right side). Common ostia were more frequent on the left side than on the right side (33% *vs.* 13%); the middle lobe drained directly into the left atrium or inferior pulmonary vein in 11% of patients; and the right inferior pulmonary vein often divided immediately at the root in 23% of patients. Fourdrain *et al.* also analyzed variations of the pulmonary arteries and veins (25,26). For pulmonary veins, 36% of patients had variations, and variations were more frequent on the right side than on the left side. The most frequent right-sided variation was the existence of three separate pulmonary veins, whereas the most frequent left-sided variation was the existence of a single pulmonary vein. Shiina *et al.* also analyzed variations of the pulmonary vein in the right upper lobe (RUL), right middle lobe (RML), right lower lobe (RLL), and left upper lobe (LUL) and reported that the incidence of variations ranged from 2.6% to 15.3%, but found no variants in the left lower lobe (LLL). They emphasized the importance of variations that could be critical during lung resection, such as anomalous V², V⁶, RML veins, and lingular veins (27).

Anatomy of the right lobes

In 2015, based on 3D-CT angiography and bronchiography

(3D-CTAB), we analyzed the anatomical variations of RUL in more detail and compared data with those in previous cadaveric studies (Table 2, Figures 2,3) (28). Although the incidence of variations in pulmonary arteries was similar, there were differences in the incidence of variations in veins and bronchi, such as the B¹⁻ or B²⁻defective patterns. Zhang *et al.* later studied the B¹⁻defective type in more detail, and additionally analyzed variations in vascular patterns (38). Based on anatomical data, we further created a simplified model of segmental anatomy to guide surgeons while performing segmentectomies (39,40). Our aim was to classify the wide-variety of segmental anatomy into several specific patterns, allowing surgeons to perform segmentectomies with a pattern-based approach. Zhang *et al.* also created a simplified anatomical model for the left upper division which can be considered to be the counterpart of the RUL (41).

Interestingly, some have studied the RUL anatomy by analyzing anatomical structures bilaterally or by taking into account lung volume as a factor for classification. Wang *et al.* compared the superior pulmonary veins bilaterally and proposed a uniform classification that could be applicable for both upper lobes, that is, classifying the veins into central, semi-central, and non-central types (22). Chen *et al.* included segmental lung volume analyzed by 3D imaging as a factor to determine the “dominant pulmonary segment” of the RUL and subsequently determined whether segmental lung volume could be correlated to anatomical variations (19). Other studies analyzed more specific features




Figure 2 Anatomical analyses of the lung based on 3D-CT imaging. Anatomical features analyzed by 3D-CT imaging are grouped according to lobes with references in parenthesis. RUL, right upper lobe; RML, right middle lobe; RLL, right lower lobe; LUL, left upper lobe; LLL, left lower lobe; PV, pulmonary vein, PA, pulmonary artery.

of the RUL such as the quadrivial pattern bronchus (42), the V²a intersegmental vein (43), the right top pulmonary vein (44), or Boyden's triad (21,23,45).

In 2017, we subsequently reported on the segmental anatomy of the RML and RLL (*Table 2, Figure 3*) (29). Pulmonary bronchi and vessels of the RML and S⁶ were classified according to the number of stems. Bronchi and pulmonary arteries of S⁷ and basal segments were classified by branching patterns. Also, the subsuperior segment (or S*), which is an independent segment between S⁶ and S¹⁰, was identified in 20.4% of cases. Studies further analyzed the subsuperior segment in more detail, not only on the right side but also bilaterally (16-18).

Anatomy of the left lobes

Between 2020 and 2022, several studies evaluated the LUL anatomy, each using slightly different classification systems (*Table 2, Figure 3*) (30-33). Isaka *et al.* also analyzed the relationship between branching patterns of bronchi, arteries, and veins, finding a significant correlation between arterial and bronchial branching patterns as well as between arterial and venous branching patterns (46). As previously mentioned, Zhang *et al.* classified the veins of the left upper division into simplified models that can be

used during segmentectomies (41). Additional studies also focused on the lingular artery, with one study suggesting that the mediastinal lingular artery might originate from the variation of B³, and that the presence of a mediastinal lingular artery also influences the venous pattern of the left upper division (47,48).

In 2020, Maki *et al.* reported the anatomy of the LLL (*Table 2, Figure 3*) (34). They also reported rare variations such as B⁷ with independent branching from the basal bronchi; subsuperior bronchus (B*); or an extrapericardial common trunk of the left pulmonary veins. Liu *et al.* proposed a classification for the mediastinal basal artery, which is a pulmonary artery that branches from the proximal part of the left pulmonary artery between the left main bronchus and the left superior pulmonary vein, proceeding directly into the lower lobe (49). The study by Maki *et al.* also included one case of a mediastinal basal artery that branched within the pericardium (34).

Interlobar vessels

Some studies have analyzed the variations of interlobar vessels (*Table 2*). Information on interlobar vessels would be important for surgeons when identifying these vessels during anatomical lung resection or when dissecting the lung

Table 2 Details of lung anatomy analyzed by 3D-CT imaging

Anatomical features	Authors (ref.)	Bronchus	Pulmonary artery	Pulmonary vein
Lung in general	Akiba <i>et al.</i> (24)	N/A	N/A	Common ostia
	Fourdrain <i>et al.</i> (25,26)	N/A	Number of branches, frequency, and variations in all lobes	Variations of venous drainage Right side: 6 types Left side: 4 types
	Shiina <i>et al.</i> (27)	N/A	N/A	Anomalous drainage of right V ² /V ³ /V ⁶ and left lingular veins
RUL	Nagashima <i>et al.</i> (28)	4 major types: trifurcated, bifurcated, defective B ¹ or B ² , quadrivial type	4 major types	4 major types
RML	Nagashima <i>et al.</i> (29)	1 type (single stem)	3 types	3 types (some drain into IPV)
RLL	Nagashima <i>et al.</i> (29)	B ⁶ /B*/B ⁷ /B ⁸⁻¹⁰ : 2/2/4/3 types	A ⁶ /A*/A ⁷ : 3/2/4 types A ⁸⁻¹⁰ : 2 major types (4 subtypes)	V ⁶ : 2 major types (some drain into SPV) V ⁸⁻¹⁰ : 2 major types (5 subtypes)
LUL	Deng <i>et al.</i> (30)	2 types (8 subtypes)	3 types (10 subtypes) of variations	2 types (5 subtypes) of variations
	Fan <i>et al.</i> (31)	LUD: 4 types	LUD: 4 types	LUD: 3 types
		Lingula: 3 types	Lingula: 3 types	Lingula: 2 types
		Uncommon variations	Uncommon variations	Uncommon variations
	Maki <i>et al.</i> (32)	LUB: 2 major types LUD: 2 major types (4 subtypes) lingula: 2 types (1 subtype) rare variations	2 types according to lingular artery (15 subtypes) Complete mediastinal lingular artery and rare variations	LUD: 3 major types (13 subtypes) Lingula: 2 major types (8 subtypes)
He <i>et al.</i> (33)	LUB: 2 types (3 subtypes) LUD: 2 types (6 subtypes) lingula: 2 types (3 subtypes)	N/A	N/A	
LLL	Maki <i>et al.</i> (34)	B ⁶ : 2 types (6 subtypes)	A ⁶ : 3 types (13 subtypes)	V ⁶ : 2 types (8 subtypes)
		Basal bronchus: 6 types	Basal artery: 5 types, rare cases	Basal vein: 8 types and lingular vein draining into the LLV
		B*, B ⁷		Common trunk, extrapericardial
Interlobar vessels	Wang <i>et al.</i> , Xu <i>et al.</i> , Murota <i>et al.</i> (35-37)	N/A	Right side: 4 types (15 subtypes) Left side: 7 types (85 subtypes)	Right side: 2 major types (30 subtypes)

For clarity, the number of types and subtypes have been simplified in some cases and the reader should refer to the original manuscript for a detailed classification. 3D, three-dimensional; CT, computed tomography; RUL, right upper lobe; RML, right middle lobe; RLL, right lower lobe; LUL, left upper lobe; LLL, left lower lobe; LUD, left upper division; LUB, left upper bronchus; PV, pulmonary vein; SPV, superior pulmonary vein; IPV, inferior pulmonary vein; LLV, left lower vein; N/A, not assessed; ref., references.




Figure 3 Anatomical classifications and subtypes of the lung based on 3D-CT images. Overview of subtypes for (A) bronchus, (B) pulmonary artery, and (C) pulmonary veins according to analyses. For clarity, the number of types and subtypes have been simplified in some cases and the reader should refer to the original manuscript for a detailed classification. RUL, right upper lobe; RML, right middle lobe; RLL, right lower lobe; LUL, left upper lobe; LLL, left lower lobe; LUB, left upper bronchus; LUD, left upper division; B*, bronchus of subsuperior segment; A*, artery of subsuperior segment.

fissure in patients with incomplete lobulation. Wang *et al.* classified the right interlobar arteries according to the order and number of branches of the RML artery and A⁶ (35). Xu *et al.* classified the general morphology of the right interlobar veins and reported that interlobar veins hidden by an incomplete upper oblique fissure were most vulnerable to accidental injury; a diameter larger than 2.4 mm for the oblique fissure interlobar vein type or less than 2 mm for the mediastinal interlobar vein type was also associated with a higher risk of injury (36). Murota *et al.* also classified left-side interlobar arteries (37).

Limitations of 3D imaging and development of 3D imaging software

Despite the advantages and contribution to our current understanding of lung anatomy as detailed above, current 3D-CT imaging has its limitations. For example, small blood vessels that are visible on conventional 2D images may not always be reconstructed in 3D-CT images. In our previous study, small vessels with a diameter of less than 1.5 mm were missed on 3D-CT images when compared to intraoperative views (28). For a better identification of small anatomical structures, observing both 3D-CT images and thin-section CT images is equally important. There will always be some inherent difference between 2D images, 3D-CT reconstructed images, and intraoperative views. Vessels that were preoperatively overlooked by 2D or 3D images might only be recognized intraoperatively. Therefore, feedback from actual intraoperative findings is important to further refine 3D-CT reconstruction methods. Use of 3D-CT images could also be limited by other factors including limited availability of 3D reconstruction software; inadequate conditions during CT examination; and the presence of tumors or atelectasis obstructing bronchi, impeding vascular flow, or obscuring peripheral anatomy. To overcome these limitations, numerous imaging software platforms have been developed and optimized, including Ziostation 2 (Ziosoft Inc.), REVORAS (Ziosoft Inc.), Synapse Vincent (Fujifilm), IQQA-Lung (EDDA Technology), Deepinsight platform (Neusoft Group Ltd.), Mimics software (Materialise), PV-iCAS (PVmed), and the list continues to grow (19,21,28,33,44,45,48). Until now, the prerequisite for 3D imaging of pulmonary vessels was the availability of contrast-enhanced CT images. However, 3D imaging software now allows for 3D reconstruction of pulmonary vessels from non-enhanced CT data (50-53).

Conclusions

We reviewed here key features of 3D-CT imaging of the lung and presented results from representative anatomical studies. Such studies have led to new or modified classification systems, shed light on lung anatomy from a useful surgical viewpoint, and enabled us to analyze lung anatomy with a focus on particular anatomical features. 3D-CT images also allow for enhanced pre- and intra-operative simulation, improved surgical safety, enhanced educational utility, and the capacity to perform large-scale anatomical studies in shorter time frames. A decade has passed since the initial reports on 3D-CT image-guided lung resection (54-59) and 3D-CT imaging has become widely implemented with results of prospective multicenter studies now being reported (60). The current surge of 3D-CT imaging analysis shows that the field is still growing, with the technology continuing to improve and now even being combined with virtual reality and artificial intelligence (61,62). Future studies using these new and innovative methodologies will continue to refine our understanding of lung anatomy while enhancing our ability to perform safe and effective surgical resections.

Acknowledgments

Funding: None.

Footnote

Provenance and Peer Review: This article was commissioned by the editorial office, *Video-Assisted Thoracic Surgery* for the series “Simulation and Navigation Techniques in VATS/RATS”. The article has undergone external peer review.

Peer Review File: Available at <https://vats.amegroups.org/article/view/10.21037/vats-23-21/prf>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://vats.amegroups.org/article/view/10.21037/vats-23-21/coif>). The series “Simulation and Navigation Techniques in VATS/RATS” was commissioned by the editorial office without any funding or sponsorship. SN served as the unpaid Guest Editor of the series. HI served as the unpaid Guest Editor of the series and serves as the unpaid Associate Editor-in-Chief of *Video-Assisted Thoracic Surgery* from December 2022 to November 2024. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. NOMENCLATURE of broncho-pulmonary anatomy; an international nomenclature accepted by the Thoracic Society. *Thorax* 1950;5:222-8.
2. Cory RA, Valentine EJ. Varying patterns of the lobar branches of the pulmonary artery. A study of 524 lungs and lobes seen at operation of 426 patients. *Thorax* 1959;14:267-80.
3. Boyden EA, Scannell JG. An analysis of variations in the bronchovascular pattern of the right upper lobe of 50 lungs. *Am J Anat* 1948;82:27-73.
4. Boyden EA, Hartmann JF. An analysis of variations in the bronchopulmonary segments of the left upper lobes of fifty lungs. *Am J Anat* 1946;79:321-60.
5. Yamashita H. Roentgenologic anatomy of the lung. *Igaku-Shoin*; 1978:389.
6. Akiba T. Utility of three-dimensional computed tomography in general thoracic surgery. *Gen Thorac Cardiovasc Surg* 2013;61:676-84.
7. Oizumi H, Endoh M, Takeda S, et al. Anatomical lung segmentectomy simulated by computed tomographic angiography. *Ann Thorac Surg* 2010;90:1382-3.
8. Oizumi H, Kanauchi N, Kato H, et al. Anatomic thoracoscopic pulmonary segmentectomy under 3-dimensional multidetector computed tomography simulation: a report of 52 consecutive cases. *J Thorac Cardiovasc Surg* 2011;141:678-82.
9. Kanzaki M, Maeda H, Wachi N, et al. Complete video-assisted thoracoscopic multi-subsegmentectomy based on patients' specific virtual 3-D pulmonary models. *Asian J Endosc Surg* 2013;6:110-5.
10. Chen-Yoshikawa TF, Date H. Update on three-dimensional image reconstruction for preoperative simulation in thoracic surgery. *J Thorac Dis* 2016;8:S295-301.
11. Shimizu K, Nakazawa S, Nagashima T, et al. 3D-CT anatomy for VATS segmentectomy. *J Vis Surg* 2017;3:88.
12. Nakazawa S, Shimizu K, Mogi A, et al. VATS segmentectomy: past, present, and future. *Gen Thorac Cardiovasc Surg* 2018;66:81-90.
13. Wang R, Zhang Y, Hu Q, et al. Identification of the segmental structures of the right upper lobe of the lung using non-enhanced thin-slice CT. *J Thorac Dis* 2020;12:1639-44.
14. Bhakhri K, Hyde ER, Mak SM, et al. Surgeon knowledge of the pulmonary arterial system and surgical plan confidence is improved by interactive virtual 3D-CT models of lung cancer patient anatomies. *Front Surg* 2021;8:652428.
15. Vervoorn MT, Wulfse M, Mohamed Hoesein FAA, et al. Application of three-dimensional computed tomography imaging and reconstructive techniques in lung surgery: A mini-review. *Front Surg* 2022;9:1079857.
16. Shimizu K, Mogi A, Yajima T, et al. Thoracoscopic subsuperior segment segmentectomy. *Ann Thorac Surg* 2017;104:e407-10.
17. Zhou D, Gao Y, Wang H, et al. Prevalence and anatomical characteristics of subsuperior segment in lung lower lobe. *J Thorac Cardiovasc Surg* 2022;S0022-5223(22)00411-1.
18. Maki R, Miyajima M, Ogura K, et al. Anatomy of the left subsuperior segment for segmentectomy. *Surg Today* 2022;52:1054-62.
19. Chen ZH, Chu XP, Zhang JT, et al. The regularity of anatomical variations of dominant pulmonary segments in the right upper lobe. *Thorac Cancer* 2023;14:462-9.
20. Mimae T, Miyata Y, Kumada T, et al. The intersegmental pulmonary vein is not always located on the intersegmental plane of the lung: Evaluation with 3-dimensional volume-rendering image reconstruction. *JTCVS Tech* 2022;16:132-8.
21. Zhang M, Sun WJ, Wu QC, et al. Boyden's triad in the left lung: an interesting phenomenon. *Interact Cardiovasc Thorac Surg* 2022;35:ivac082.
22. Wang J, Lin H, Bian C, et al. A modified system for classifying the bilateral superior pulmonary veins using three-dimensional computed tomography bronchography and angiography images. *J Thorac Dis* 2021;13:5933-41.
23. Zhang M, Mao N, Wu Q, et al. Boyden's triad: the past, present and future. *Interact Cardiovasc Thorac Surg* 2022;34:590-6.

24. Akiba T, Marushima H, Odaka M, et al. Pulmonary vein analysis using three-dimensional computed tomography angiography for thoracic surgery. *Gen Thorac Cardiovasc Surg* 2010;58:331-5.
25. Fourdrain A, De Dominicis F, Bensussan M, et al. Three-dimensional computed tomography angiography of the pulmonary veins and their anatomical variations: involvement in video-assisted thoracoscopic surgery-lobectomy for lung cancer. *Folia Morphol (Warsz)* 2017;76:388-93.
26. Fourdrain A, De Dominicis F, Blanchard C, et al. Three-dimensional CT angiography of anatomic variations in the pulmonary arterial tree. *Surg Radiol Anat* 2018;40:45-53.
27. Shiina N, Kaga K, Hida Y, et al. Variations of pulmonary vein drainage critical for lung resection assessed by three-dimensional computed tomography angiography. *Thorac Cancer* 2018;9:584-8.
28. Nagashima T, Shimizu K, Ohtaki Y, et al. An analysis of variations in the bronchovascular pattern of the right upper lobe using three-dimensional CT angiography and bronchography. *Gen Thorac Cardiovasc Surg* 2015;63:354-60.
29. Nagashima T, Shimizu K, Ohtaki Y, et al. Analysis of variation in bronchovascular pattern of the right middle and lower lobes of the lung using three-dimensional CT angiography and bronchography. *Gen Thorac Cardiovasc Surg* 2017;65:343-9.
30. Deng Y, Cai S, Huang C, et al. Anatomical variation analysis of left upper pulmonary blood vessels and bronchi based on three-dimensional reconstruction of chest CT. *Front Oncol* 2022;12:1028467.
31. Fan K, Feng J, Li Y, et al. Application of three-dimensional reconstruction of left upper lung lobes in anatomical segmental resection. *Thorac Cancer* 2022;13:1176-83.
32. Maki R, Miyajima M, Ogura K, et al. Pulmonary vessels and bronchus anatomy of the left upper lobe. *Surg Today* 2022;52:550-8.
33. He H, Wang F, Wang PY, et al. Anatomical analysis of variations in the bronchus pattern of the left upper lobe using three-dimensional computed tomography angiography and bronchography. *Ann Transl Med* 2022;10:305.
34. Maki R, Miyajima M, Ogura K, et al. Pulmonary vessels and bronchial anatomy of the left lower lobe. *Surg Today* 2020;50:1081-90.
35. Wang LF, Zhao L, Lv CS, et al. Anatomical type analysis of right interlobar artery based on chest thin-slice CT scan and three-dimensional reconstruction. *J Cardiothorac Surg* 2022;17:328.
36. Xu X, Wang J, Liu Q, et al. Clinical significance of the diverse interlobar veins hidden in the upper oblique fissure. *Ann Transl Med* 2022;10:34.
37. Murota M, Yamamoto Y, Satoh K, et al. An analysis of anatomical variations of the left pulmonary artery of the interlobar portion for lung resection by three-dimensional CT pulmonary angiography and thin-section images. *Jpn J Radiol* 2020;38:1158-68.
38. Zhang M, Mao N, Wang SH, et al. The B1 defective type of bifurcated right upper lobe bronchus. *J Thorac Dis* 2019;11:4218-23.
39. Shimizu K, Nagashima T, Ohtaki Y, et al. Analysis of the variation pattern in right upper pulmonary veins and establishment of simplified vein models for anatomical segmentectomy. *Gen Thorac Cardiovasc Surg* 2016;64:604-11.
40. Nakazawa S, Shimizu K, Kawatani N, et al. Right upper lobe segmentectomy guided by simplified anatomic models. *JTCVS Tech* 2020;4:288-97.
41. Zhang M, Mao N, Zhang K, et al. Analysis of the variation pattern in left upper division veins and establishment of simplified vein models for anatomical segmentectomy. *Ann Transl Med* 2020;8:1515.
42. Zhong X, Huang Y, Yang M, et al. Elucidating the anatomy of the quadrivial pattern of the right upper lobe bronchus using 3D-CT images. *Ann Transl Med* 2022;10:46.
43. Zhou H, Wei W, He H, et al. A cross-sectional study: analysis of anatomical variation in the right upper lung intersegmental vein V2a based on a 3D reconstruction technique. *J Thorac Dis* 2022;14:4460-7.
44. Miyamoto N, Yoshida M, Takashima M, et al. Classifying the destination of right top pulmonary vein in 31 clinical cases. *Gen Thorac Cardiovasc Surg* 2021;69:1192-5.
45. Miura K, Shimizu K, Mishima S, et al. Anatomical resection for right B(3) downwards-shifting malformation. *Gen Thorac Cardiovasc Surg* 2023;71:71-5.
46. Isaka T, Mitsuboshi S, Maeda H, et al. Anatomical analysis of the left upper lobe of lung on three-dimensional images with focusing the branching pattern of the subsegmental veins. *J Cardiothorac Surg* 2020;15:273.
47. He H, Chen P, Chen X, et al. Analysis of anatomical variations of the lingular artery of the left upper lobe using 3D computed tomography angiography and bronchography. *J Thorac Dis* 2021;13:5035-41.
48. Gao C, Xu WZ, Li ZH, et al. Analysis of bronchial and vascular patterns in left upper lobes to explore the genesis of mediastinal lingular artery and its influence on pulmonary

- anatomical variation. *J Cardiothorac Surg* 2021;16:306.
49. Liu Y, Zhang S. Mediastinal basal pulmonary artery identification and classification by three-dimensional reconstruction. *Surg Radiol Anat* 2022;44:447-53.
 50. Chen X, Wang Z, Qi Q, et al. A fully automated noncontrast CT 3-D reconstruction algorithm enabled accurate anatomical demonstration for lung segmentectomy. *Thorac Cancer* 2022;13:795-803.
 51. Hanawa R, Shigenobu T, Tajima A. Three-dimensional reconstruction of pulmonary blood vasculature from plain computed tomography using Ziostation2. *Jpn J Chest Surg* 2019;33:578-86.
 52. Nakao M, Omura K, Hashimoto K, et al. Three-dimensional image simulation for lung segmentectomy from unenhanced computed tomography data. *Gen Thorac Cardiovasc Surg* 2022;70:312-4.
 53. Nakazawa S, Hanawa R, Nagashima T, et al. Segmentectomy guided by 3-dimensional images reconstructed from nonenhanced computed tomographic data. *Ann Thorac Surg* 2021;111:e301-4.
 54. Shimizu K, Nakano T, Kamiyoshihara M, et al. Segmentectomy guided by three-dimensional computed tomography angiography and bronchography. *Interact Cardiovasc Thorac Surg* 2012;15:194-6.
 55. Iwano S, Usami N, Yokoi K, et al. Segmentectomy simulation using a virtual three-dimensional safety margin. *Ann Thorac Surg* 2012;93:e37-9.
 56. Nakano T, Shimizu K, Nakano S, et al. Usefulness of three-dimensional computed tomographic angiography with bronchography for the planning of minimally invasive video-assisted thoracic surgery for intralobar pulmonary sequestration. *Eur J Cardiothorac Surg* 2013;43:199.
 57. Ikeda N, Yoshimura A, Hagiwara M, et al. Three dimensional computed tomography lung modeling is useful in simulation and navigation of lung cancer surgery. *Ann Thorac Cardiovasc Surg* 2013;19:1-5.
 58. Saji H, Inoue T, Kato Y, et al. Virtual segmentectomy based on high-quality three-dimensional lung modelling from computed tomography images. *Interact Cardiovasc Thorac Surg* 2013;17:227-32.
 59. Iwano S, Yokoi K, Taniguchi T, et al. Planning of segmentectomy using three-dimensional computed tomography angiography with a virtual safety margin: technique and initial experience. *Lung Cancer* 2013;81:410-5.
 60. Niu Z, Chen K, Jin R, et al. Three-dimensional computed tomography reconstruction in video-assisted thoracoscopic segmentectomy (DRIVATS): A prospective, multicenter randomized controlled trial. *Front Surg* 2022;9:941582.
 61. Sadeghi AH, Maat APWM, Taverne YJHJ, et al. Virtual reality and artificial intelligence for 3-dimensional planning of lung segmentectomies. *JTCVS Tech* 2021;7:309-21.
 62. Bakhuis W, Kersten CM, Sadeghi AH, et al. Preoperative visualization of congenital lung abnormalities: hybridizing artificial intelligence and virtual reality. *Eur J Cardiothorac Surg* 2022;63:ezad014.

doi: 10.21037/vats-23-21

Cite this article as: Nakazawa S, Nagashima T, Kawatani N, Gedeon PC, DeSimone AK, Igai H, Kosaka T, Shirabe K. Anatomy of the lung revisited by 3D-CT imaging. *Video-assist Thorac Surg* 2023;8:17.