

Supplementary




Figure S1 Kaplan-Meier analysis of progression-free survival (PFS) according to interval time. Kaplan-Meier curves comparing PFS between patients with an interval time ≤ 10 months and those with an interval time > 10 months. PFS, progression-free survival.




Figure S2 Kaplan-Meier analysis of overall survival (OS) according to interval time. Kaplan-Meier curves comparing OS between patients with an interval time ≤ 10 months and those with an interval time > 10 months. OS, overall survival.




Figure S3 Kaplan-Meier analysis of progression-free survival (PFS) according to T790M mutation status. Kaplan-Meier curves showing PFS in patients with T790M-negative status versus those with T790M-positive status. PFS, progression-free survival.




Figure S4 Kaplan-Meier analysis of overall survival (OS) according to T790M mutation status. Kaplan-Meier curves showing OS in patients with T790M-negative status versus those with T790M-positive status. OS, overall survival.




Figure S5 Kaplan-Meier analysis of progression-free survival (PFS) according to TP53 mutation status. Kaplan-Meier curves comparing PFS between patients with TP53 wild-type and those with TP53 mutation. PFS, progression-free survival.




Figure S6 Kaplan-Meier analysis of overall survival (OS) according to TP53 mutation status. Kaplan-Meier curves comparing OS between patients with TP53 wild-type and those with TP53 mutation. OS, overall survival.