

Reference *Figure 5*

1. Haydock D, Barratt-Boyes B, Macedo T, et al. Aortic valve replacement for active infectious endocarditis in 108 patients. A comparison of freehand allograft valves with mechanical prostheses and bioprostheses. *J Thorac Cardiovasc Surg* 1992;103:130-9.
2. Kaiser SP, Melby SJ, Zierer A, et al. Long-term outcomes in valve replacement surgery for infective endocarditis. *Ann Thorac Surg* 2007;83:30-5.
3. Lytle BW, Priest BP, Taylor PC, et al. Surgical treatment of prosthetic valve endocarditis. *J Thorac Cardiovasc Surg* 1996; 111:198-207; discussion 10.
4. Grinda JM, Mainardi JL, D'Attellis N, et al. Cryopreserved aortic viable homograft for active aortic endocarditis. *Ann Thorac Surg* 2005;79:767-71.
5. Perrotta S, Lentini S. In patients with severe active aortic valve endocarditis, is a stentless valve as good as the homograft ? *Interact Cardiovasc Thorac Surg* 2010;11:309-13.
6. Flameng W, Daenen W, Jashari R, et al. Durability of homografts used to treat complex aortic valve endocarditis. *Ann Thorac Surg* 2015;99:1234-8.
7. David TE, Feindel CM, Bos J, et al. Aortic valve replacement with Toronto SPV bioprosthesis: optimal patient survival but suboptimal valve durability. *J Thorac Cardiovasc Surg* 2008;135:19-24.
8. Ozaki S, Kawase I, Yamashita H, et al. A total of 404 cases of aortic valve reconstruction with glutaraldehyde-treated autologous pericardium. *J Thorac Cardiovasc Surg* 2014;147:301-6.