

**Table S1** First-line and second line treatment modalities in 124 patients (133 breast)

Treatment modalities	Number (%)
First-line treatment after breast clinic evaluation	
Antibiotic <sup>†</sup>	14 (10.5)
Steroids <sup>‡</sup>	56 (42.1)
Antibiotics + steroids	12 (9.0)
Surgical excision or I/D <sup>§</sup>	7 (5.3)
Observation	41 (30.8)
Other <sup>¶</sup>	3 (2.3)
Second line treatment after first follow-up	
Antibiotics <sup>‡</sup>	9 (6.8)
Methotrexate	1 (0.8)
Observation	44 (33.1)
Steroids	32 (24.1)
Surgical excision or I/D <sup>*</sup>	5 (3.8)
No follow-up ultrasound	42 (31.6)
Total	133 (100.0)

<sup>†</sup>, 4/14 (28.6%) were treated with both antibiotics and incision and drainage; <sup>‡</sup>, 4/56 (7.1%) were treated only with topical steroids; <sup>§</sup>, 2/7 (28.6%) were only treated with incision and drainage. 1 of the remaining 5 patients (1/5, 20%) was only treated with surgical excision. 3/5 (60.0%) were treated with surgical excision and steroids. Likewise, 1/5 (20%) was also treated with antibiotics in addition to surgical excision; <sup>¶</sup>, other includes, two patients who were lost to follow-up and one patient who was treated with antibiotics, steroids, and incision and drainage; <sup>‡</sup>, 7/9 used steroids in addition to antibiotics; <sup>\*</sup>, 1/5 was treated with only incision and drainage. I/D, incision and drainage.