



**Figure S1** Risk of bias across studies: funnel plot of studies included in our meta-analysis of prevalence of Charles Bonnet syndrome in low vision patients.

**Table S1** Sensitivity analysis on the estimate of the prevalence of Charles Bonnet syndrome in low vision patients

Excluded study	Pooled prevalence	LCI 95%	HCI 95%	Cochran Q	I <sup>2</sup>
Shiraishi <i>et al.</i> 2004	22.5%	16.8%	28.8%	134.3	93.3
Elflein <i>et al.</i> 2016	21.3%	15.0%	28.3%	178.1	94.9
Teunisse <i>et al.</i> 1995	20.5%	14.0%	27.9%	184.5	95.1
Marmamula <i>et al.</i> 2019	20.5%	13.3%	28.8%	175.3	94.9
Tatlipinar <i>et al.</i> 2001	20.2%	13.9%	27.3%	188.4	95.2
Vikucevic & Fitzmaurice 2008	20.0%	13.5%	27.4%	188.8	95.2
Gordon 2016	19.9%	11.8%	29.4%	187.9	95.2
Crumbly <i>et al.</i> 2008	19.4%	13.2%	26.3%	187.7	95.2
Gilmour <i>et al.</i> 2009	18.4%	12.6%	24.9%	154.0	94.2
Jackson <i>et al.</i> 2007	17.8%	12.3%	24.2%	155.6	94.2
Menon 2005	16.9%	11.7%	22.7%	146.1	93.8

LCI 95%, lower 95% confidence interval; HCI, higher/upper 95% confidence interval.