

**Table S1** Results of the dynamic risk stratification at 12 months and at the end of follow-up in patients with available data


Type of response	12 months (n=148)		Last visit (n=164)	
	n	%	n	%
Excellent	106	71.6	122	74.4
Indeterminate	36	24.3	34	20.7
Biochemical incomplete	5	3.4	6	3.7
Structural incomplete	1	0.7	2	1.2

n, number of subjects.


**Table S2** Comparison of 148 patients at the end of the first year of follow-up classified according to absence (excellent response) or presence of disease (any other type of response)

Characteristics	n	Excellent response (n=106)	Any other response (n=42)	P
Age, years	148	49.5±4.4	54.9±14.3	0.044
Duration of follow-up, months	148	45.4 (22.3–65.9)	44.3 (23.1–66.7)	0.973
Gender	148			0.655
Female		85 (80.2)	32 (76.2)	
Male		21 (19.8)	10 (23.8)	
Histological type	148			1.000
Papillary		102 (96.2)	40 (95.2)	
Follicular		4 (3.8)	2 (4.8)	
Tumor size, cm	148	0.4 (0.24–0.7)	0.4 (0.2–0.63)	0.792
Chronic lymphocytic thyroiditis	148	19 (17.9)	20 (47.6)	<0.001
Multifocality	148			1.000
Unifocal		102 (96.2)	40 (95.2)	
Multifocal		4 (3.8)	2 (4.8)	
Incidental	148	78 (73.6)	34 (81.0)	0.284
Central lymph node dissection	147	3 (2.9)	2 (4.8)	0.624
Risk of recurrence ATA 2015	148			0.578
Low		102 (96.2)	42 (100.0)	
Intermediate		4 (3.8)	0 (0)	
Nodules at remnant lobe	148	26 (24.5)	20 (47.6)	0.010
Size of the largest nodule in remnant lobe, cm	39	1.0 (0.6–1.3)	0.9 (0.6–1.3)	0.813

Data are the mean ± SD or the median (interquartile range) for quantitative variables, and the number (percentage) for categorical variables. n, number of subjects; ATA, American Thyroid Association.



**Figure S1** Diagram indicating the criteria for indeterminate response at the end of follow-up in patients with papillary and follicular thyroid carcinoma. TgAb, thyroglobulin antibodies.



**Figure S2** Diagram indicating the evolution of patients classified according to the presence or absence of nodules during follow-up. The reason for completing the thyroidectomy in the patient with no nodules during follow-up was the presence of suspicious cervical lymph nodes. PTC, papillary thyroid carcinoma; FTC, follicular thyroid carcinoma.