

Table S1 Antimicrobial susceptibility of the isolated *Klebsiella pneumoniae*

Antibacterials	Strain A	Strain B	Strain C	Strain D
	Blood sample	Pus sample	Blood sample	BALF sample
Amikacin	≤2 (S)	≤2 (S)	≥64 (R)	≥64 (R)
Ceftazidime	≤1 (S)	≤1 (S)	≥64 (R)	≥64 (R)
Ceftriaxone	≤1 (S)	≤1 (S)	≥64 (R)	≥64 (R)
Cefepime	≤1 (S)	≤1 (S)	≥64 (R)	≥64 (R)
Imipenem	≤1 (S)	≤1 (S)	≥16 (R)	≥16 (R)
Ampicillin-Sulbactam	4 (S)	4 (S)	≥32 (R)	≥32 (R)
Tobramycin	≤1 (S)	≤1 (S)	≥16 (R)	≥16 (R)
Aztreonam	≤1 (S)	≤1 (S)	≥64 (R)	≥64 (R)
Ciprofloxacin	≤0.25 (S)	≤0.25 (S)	≥4 (R)	≥4 (R)
Cefotetan	≤4 (S)	≤4 (S)	≥64 (R)	≥64 (R)
Ertapenem	≤0.5 (S)	≤0.5 (S)	≥16 (R)	≥16 (R)
Gentamycin	≤1 (S)	≤1 (S)	≥16 (R)	≥16 (R)
Levofloxacin	≤0.25 (S)	≤0.25 (S)	≥8 (R)	≥8 (R)
Sulfamethoxazole-Trimethoprim	≤20 (S)	≤20 (S)	≥320 (R)	≥320 (R)
Piperacillin-Tazobactam	≤4 (S)	≤4 (S)	≥128 (R)	≥128 (R)

Detection method: minimal inhibitory concentrations (ug/mL). S, Susceptibility; R, Resistance.