

Table S1 The important test results of the patient after admission

Test	Result	Range of reference values
Coagulation tests		
PT, s	11.4	9.8–12.1
PTA, %	107.7	70.0–130.0
PT/R	1	0.82–1.15
PTINR	1	0.82–1.15
Fbg, g/L	3.05	2–4
APTT, s	25.8	25.0–31.3
TT, s	17.5	14.0–21.0
D-dimer, mg/L FEU	0.18	0.00–0.55
PIVKA-II, mAU/mL	27	0–40
Digestive tumor series		
AFP, ng/mL	7.5↑	≤7
CEA, ng/mL	1.5	≤5.0
CA125, U/mL	3.4	≤24
CA19-9, U/mL	14	≤30
CA72-4, U/mL	2.2	≤6.9
Blood-cell analysis		
WBC count, 10 ⁹ /L	5.23	3.5–9.5
RBC count, 10 ¹² /L	4.68	4.30–5.80
HGB count, g/L	139	130–175
PLT count, 10 ⁹ /L	95↓	125–350
NEUT%, %	75.7↑	45–75
Biochemical tests		
ALT, U/L	20	9–50
AST, U/L	35	15–40
TBIL, μmol/L	12.4	≤26.0
DBIL, μmol/L	3	0.0–3.4
IBIL, μmol/L	9.4	3.4–18.8
GGT, U/L	71	10–60
ALP, U/L	66	45–125
ALB, g/L	46.8	40–55
Urea, mmol/L	10.5	3.6–9.5
Crea, μmol/L	101	57–111
Bun/Crea	25.7↑	12–20

Table S1 (continued)

Table S1 (*continued*)

Test	Result	Range of reference values
Hepatitis B virus tests		
HBsAg, IU/mL	>250.00↑	<0.08
HBsAb, IU/L	<2.00	<10
HBeAg, COI	0.17	>1.00
Anti-HBe, COI	0.07↓	>1.00
Anti-HBc, COI	0.00↓	>1.00
HBV-DNA, IU/mL	<1.00×10 ²	<1.00×10 ²

PT, prothrombin time; PTA, prothrombin time activity; PT/R, prothrombin time/ratio; PTINR, prothrombin time international normalized ratio; Fbg, fibrinogen; APTT, activated partial thromboplastin time; TT, thrombin time; PIVKA-II, protein induced by vitamin K absence or antagonist-II; AFP, alpha-fetoprotein; CEA, carcinoembryonic antigen; CA125, carbohydrate antigen 125; CA19-9, carbohydrate antigen 19-9; CA72-4, carbohydrate antigen 72-4; WBC, white blood cell; RBC, red blood cell; HGB, hemoglobin; PLT, platelet; NEUT%, neutrophil%; ALT, alanine transaminase; AST, aspartate transaminase; TBIL, total bilirubin; DBIL, direct bilirubin; IBIL, indirect bilirubin; GGT, γ-glutamyl transpeptidase; ALP, alkaline phosphatase; ALB, albumin; Urea, blood urea nitrogen; Crea, Creatinine; Bun/Crea, blood urea nitrogen/creatinine.