Table S1 Reported cases of PACC received anti-tumor drug

Reported cases of PACC received anti-	-tumor drug								
DOI (ref.)	Author		Age (years)	. ,	Altered gene				Clinical course
10.1007/s10689-024-00390-3 (11)	Matsubayashi	2024	73	М	BRCA2	Liver metastasis in 13 months after surgery	olaparib	PR	N/A
10.1007/s00432-024-05841-z (12)	Merz	2024	69	М	MLH1/MSI-high	Locally advanced	mFFX	SD	SD with mFFX in 6 cycles. Sent for GnP (PD). PR with pembrolizumab. Sent for conversion surgery
10.1002/cnr2.70007 (13)	Kubo	2024	53	М	BRCA1	Liver metastasis after surgery	GnP	PD	PR with mFFX in 19 cycles. Switch to olaparib but PD in 4 months. PD with mFFX. PD with GEM
10.1007/s12328-024-01981-4 (4)	Urabe	2024	80	М	BRCA2	Relapse in remnant pancreas after	GnP	PD	PD with S1 in 8 months. Radiation therapy. PD with S1 in 5 months. PR with mFFX (continued a year
10.1007/s12328-024-01992-1 (30)	Watanabe	2024	60	М	BRCA1	surgery  Locally advanced	GnP	PD	3 months). Switch to olaparib (maintained PR in a year). Survival in a year after stopping olaparib PR with FFX (6 cycles). Sent for conversion surgery. FFX as adjuvant chemotherapy in 6 months. N
10.1002/gcc.23222 (38)	Von Fritsch	2024	52	М	BRAFV600E	Peritoneal dissemination	mFFX	SD	recurrence in 10 months after stopping FFX  SD with mFFX in 8 cycles. Stop by peripheral neurological disorders and fatigue. CR with dabrafer trametinib in 16 months
10 12008/wice v11 i2/ 5823 /20\	Wana	2023	77	М	ROS1-CENPW	Liver metastasis	Dabrafenib/trametinib	PR	N/A
10.12998/wjcc.v11.i24.5823 (39) 10.3389/fonc.2024.1357233 (24)	Wang Wu	2023	45	IVI	High TMB	Locally advanced	Tripalimab		N/A
10.3748/wjg.v28.i45.6421 (33)	Lee	2024	70	M	BRCA2	Liver metastasis	mFFX		PR with mFFX in 15 cycles. Switch to olaparib due to peripheral neurological disorders. Maintainer
10.1002/cnr2.1648 (3)	Uemura	2022	67	M	N/A	Peritoneal dissemination	FFX		5 months  PR with FFX in 19 months. Sent for conversion surgery. Adjuvant therapy with S1. No recurrence in
	00111010		0.						months after surgery
10.1002/ccr3.6718 (34)	Lelong	2022	38	М	BRCA2	Liver metastasis	FFX	PR	PR with FFX in 12 cycles. Switch to olaparib (maintain PR in 24 months)
10.1007/s12328-021-01571-8 (25)	Izumo	2022	65	М	N/A	Locally advanced (portal thrombus)	mFFX	PR	PR with mFFX in 14 cycles. Sent for conversion surgery. No recurrence in 33 months after surgery
10.1007/s12328-021-01518-z (2)	Maehira	2021	48	F	N/A	Liver metastasis after surgery	Cisplatin plus irrinotecan	PR	PR with cisplatin plus IRI in 7 months. Switch to conversion surgery. No relapse in 33 months after
10.6004/jnccn.2020.7641 (40)	Gupta	2021	81	М	NTRK fusion gene	Lymph-node relapse after surgery	GEM	ΡD	PD with GEM in 5 cycles. PD with GnP in 4 cycles
10.1200/PO.21.00400 (21)	Gaule	2022	62	M	Rearranged ALK	Liver metastasis after surgery	GnP		PD with GnP in 3 cycles. PR with nal-IRI but PD in 6 months. PD with FOLFOX in 7 cycles. PR with Alectinib in 16 months
10.1177/0300891620980792 (41)	Xu	2020	68	М	PD-L1 positive plus	Lung, retroperitoneum, and lymph- node metastasis	GnP	SD	SD with GnP in 2cycles and skin rash. PD with SOX in 2 cycles. PR with SOX plus tripalimab
10.1159/000515267 (22)	Dreikhausen	2021	66	М	BRCA2	Liver metastasis	FFX	PR	PR with FFX. Exchange to FORFIRI by peripheral neurological disorders. Sent for conversion surger FORFIRI, GnP, and nal-IRI had PD for relapse in the liver. Olaparib was sent but the efficacy was n
10.3389/fonc.2021.692480 (42)	Qin	2021	48	F	PD-L1 positive plus	Liver metastasis	Sintilimab plus	PR	available PR with sintilimab plus lenvatinib in 21 months
10 1007/-10000 000 011 (0 /07)	A BB	0000	7.1		high TMB	Liver metastasis	lenvatinib	-	DD with FFV in 6 avalog Cont for conversion and the control of the
10.1007/s12328-020-01146-z (26)	Villano	2020	71	M		Liver metastasis	FFX FOLFOX		PR with FFX in 6 cycles. Sent for conversion surgery. No relapse in 6 months after surgery
10.1016/j.radcr.2019.11.005 (46) 10.1080/15384047.2019.1595274	Fontent Kryklyva	2019 2019	56 52	M M	N/A BRCA2	Liver metastasis  Metastasis for Liver, lung.	FOLFOX FFX		PD with FOLFOX in 14 months. Sent for surgery  PR with FFX in 24 months. Stopped FFX due to peripheral neurological disorders. Death in 3 months.
(18)	,,		-			peritoneum, and skin in 6 months after surgery			
10.2169/internalmedicine.0294-17 (47)	Yoshida	2018	79	F	N/A	Liver metastasis	S1	PR	PR with S1 in 3 years
10.4251/wjgo.v10.i4.103 (43)	Richard	2018	54	М	EGFR	Liver metastasis	FFX	PD	PD with FFX in 4 cycles. PD with GnP in 2 cycles. PR with panitumumab
10.1097/MD.000000000013113 (35)	Li	2018	59	М	BRCA2	Retroperitoneum lymph node metastasis	FFX		Radiation for lymph-node metastasis (not effective). Intra-tumoral brachytherapy by radioiodine-effective). PR with olaparib. Stopped olaparib due to severe complications. PD and death
N/A (48)	Kanematsu	2018	55	F	N/A	Liver and lung metastasis after surgery	GEM	PD	TAE (cisplatin plus 5-FU) plus microwave coagulation therapy and GEM
10.1097/MD.0000000000007904 (49)	Luo	2017	41	М	N/A	Locally advanced (portal thrombus)	FFX	SD	SD with FFX in 4 cycles. Survival in 19 months
10.1007/s12328-017-0785-5 (14)	Hashimoto	2017	67	М	N/A	-	mFFX	PR	PR with mFFX in 9 months
10.1159/000445867 (17)	Bechade	2016	59	N/A	N/A	Liver metastasis	GEM plus oxaliplatin	PR	PR with GEM plus oxaliplatin in 6 cycles. Radiofrequency ablation for liver tumor. No relapse in 6
10.1089/crpc.2016.0009 (23)	Jimbo	2016	56	М	N/A	Locally advanced (CA)	FFX	SD	SD with FFX in 3 cycles. PR with FOLFOX-6 in 6 cycles. Sent for conversion surgery
10.1080/17843286.2016.1168065 (36)	Naeyaert	2016	59	М	BRCA2	Liver metastasis	GEM		PD with GEM. SD with carboplatin
N/A (27)	Toda	2016	52	M	N/A	Liver metastasis	GEM plus S1		PR with GEM plus S1 in 5 cycles. Chemo-radiation therapy and surgery. No relapse in 59 months
10.1159/000441414 (50)	Ploquin	2015	50	М	N/A	Liver metastasis	GEM plus oxaliplatin	CR	CR with GEM plus oxaliplatin in 9 cycles. Stop chemotherapy by peripheral neurological disorder relapse in 14 years
10.1016/j.pan.2015.04.006 (51)	Callata-	2015	36	F	N/A	Liver metastasis, peritoneal dissemination	Capecitabine plus temozolomide	PD	PD with capecitabine plus temozolomide in 2 cycles. SD with FFX in 9 months and PD in 11 mon with GnP
10.1016/j.pan.2015.09.006 (28)	Carhuapoma Jaunch	2016	61	F	N/A	Liver metastasis	XEROX	PR	PR with XEROX in 11 cycles. Sent for conversion surgery. Relapse at liver in 30 months after surge with XEROX in 3 cycles. No relapse with capecitabine in 9 months
10.1055/s-0033-1356439 (52)	Schempf	2014	63	М	N/A	Liver metastasis	FFX	PR	PR with FFX in 10 cycles. Stop chemotherapy in 6 weeks. PR with FORFIRI in 7 months. Maintain after stopping FORFIRI in 6 months
10.1177/030089161309900230	Ang	2013	71	М	N/A	Liver metastasis	GEM plus cisplatin	PR	PR with GEM plus cisplatin in 6 months. Renal failure. PD with GEM plus oxaliplatin. PR with FOI a year. PR with HAI (floxuridine) plus IRI. IRI plus cetuximub in 1 year. PD with FORFIRI plus cetu
(44)			25	F	N/A	Liver metastasis, peritoneal dissemination after excision of	GEM	PD	PD with GEM. Death in 2 months
(44) 10.2169/internalmedicine.51.6536 (53)	Hiraoka	2012	25			and a construct the construction			
10.2169/internalmedicine.51.6536 (53) 10.3109/0284186X.2011.617388	Hiraoka Simon	2012	79	М	N/A	primary and hepatic lesion Liver metastasis	GEM	SD	PD with GEM in 2 months. PR with FOLFOX-4 in 12 cycles. Stop chemotherapy by peripheral neurological disorders and relapse. PR with FOLFOX in 12 cycles
10.2169/internalmedicine.51.6536 (53) 10.3109/0284186X.2011.617388 (54)				M M	N/A N/A				neurological disorders and relapse. PR with FOLFOX in 12 cycles
10.2169/internalmedicine.51.6536 (53) 10.3109/0284186X.2011.617388	Simon	2011	79			Liver metastasis	GEM S1 GEM plus S1	CR	
10.2169/internalmedicine.51.6536 (53) 10.3109/0284186X.2011.617388 (54) N/A (29)	Simon	2011 2012	79 64	М	N/A	Liver metastasis  Peritoneal dissemination	S1	CR PR	neurological disorders and relapse. PR with FOLFOX in 12 cycles CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)	Simon Yamamoto Nishimizu	2011 2012 2011	79 64 64	M M	N/A N/A	Liver metastasis  Peritoneal dissemination  Liver metastasis	S1 GEM plus S1	CR PR CR	neurological disorders and relapse. PR with FOLFOX in 12 cycles  CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years  PR with GEM plus S1 in 6 cycles  CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)  N/A (56)	Simon  Yamamoto  Nishimizu  Morishima	2011 2012 2011 2010	79 64 64 65	М М М	N/A N/A N/A	Liver metastasis  Peritoneal dissemination  Liver metastasis  Liver metastasis after surgery	S1 GEM plus S1 S1	CR PR CR PD	neurological disorders and relapse. PR with FOLFOX in 12 cycles  CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years  PR with GEM plus S1 in 6 cycles  CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)  N/A (56)  N/A (57)  10.1111/j.1365-2354.2008.01014.	Simon  Yamamoto  Nishimizu  Morishima  Fujii	2011 2012 2011 2010 2010	79 64 64 65 59	М М М	N/A N/A N/A N/A	Liver metastasis  Peritoneal dissemination  Liver metastasis  Liver metastasis after surgery  Liver metastasis after surgery  Liver metastasis after surgery	S1 GEM plus S1 S1 GEM	CR PR CR PD	neurological disorders and relapse. PR with FOLFOX in 12 cycles CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years PR with GEM plus S1 in 6 cycles CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery PD with GEM in 3 cycles. PD with S1 in 3 cycles. PR with HAI (cisplatin). Relapse in 20 months at
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)  N/A (56)  N/A (57)  10.1111/j.1365-2354.2008.01014. x (58)	Simon  Yamamoto Nishimizu  Morishima  Fujii  Sorscher	2011 2012 2011 2010 2010 2009	79 64 64 65 59 48	M M M M	N/A N/A N/A N/A	Liver metastasis  Peritoneal dissemination  Liver metastasis  Liver metastasis after surgery  Liver metastasis after surgery  Liver metastasis after excision of primary and hepatic lesion  Locally advanced (portal vein)  Liver metastasis after excision of	S1 GEM plus S1 S1 GEM Paclitaxel	CR PR CR PD PD	neurological disorders and relapse. PR with FOLFOX in 12 cycles  CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years  PR with GEM plus S1 in 6 cycles  CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery  PD with GEM in 3 cycles. PD with S1 in 3 cycles. PR with HAI (cisplatin). Relapse in 20 months a  PD with weekly paclitaxel in 6 cycles. PR with GEM plus 5-FU in 9 cycles  PR with 5-FU in a year. Sent for conversion surgery. No relapse in 18 months after surgery  SD with GEM in 6months. Excision of new liver metastatic lesion. Relapse in remnant liver in 7 m
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)  N/A (56)  N/A (57)  10.1111/j.1365-2354.2008.01014. x (58)  10.1186/1477-7819-7-22 (16)	Simon  Yamamoto Nishimizu Morishima Fujii Sorscher  Distler	2011 2012 2011 2010 2010 2009	79 64 64 65 59 48	М М М М	N/A N/A N/A N/A N/A	Liver metastasis  Peritoneal dissemination  Liver metastasis  Liver metastasis after surgery  Liver metastasis after surgery  Liver metastasis after excision of primary and hepatic lesion  Locally advanced (portal vein)  Liver metastasis after excision of primary and hepatic lesion  Liver, bone, and lymph-nodes	S1 GEM plus S1 S1 GEM Paclitaxel 5-FU	CR PR CR PD PD	neurological disorders and relapse. PR with FOLFOX in 12 cycles  CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years  PR with GEM plus S1 in 6 cycles  CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery  PD with GEM in 3 cycles. PD with S1 in 3 cycles. PR with HAI (cisplatin). Relapse in 20 months a  PD with weekly paclitaxel in 6 cycles. PR with GEM plus 5-FU in 9 cycles  PR with 5-FU in a year. Sent for conversion surgery. No relapse in 18 months after surgery
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)  N/A (56)  N/A (57)  10.1111/j.1365-2354.2008.01014.  x (58)  10.1186/1477-7819-7-22 (16)  10.1093/jjco/hyp085 (59)	Simon  Yamamoto Nishimizu Morishima Fujii Sorscher  Distler Seki Seki	2011 2012 2011 2010 2010 2009 2009 2009	79 64 64 65 59 48 65 48	M M M M M	N/A N/A N/A N/A N/A N/A	Liver metastasis  Peritoneal dissemination Liver metastasis Liver metastasis after surgery Liver metastasis after surgery Liver metastasis after excision of primary and hepatic lesion Locally advanced (portal vein) Liver metastasis after excision of primary and hepatic lesion Liver, bone, and lymph-nodes metastasis	S1 GEM plus S1 S1 GEM Paclitaxel 5-FU GEM GEM	CR PR CR PD PD PD	neurological disorders and relapse. PR with FOLFOX in 12 cycles  CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years  PR with GEM plus S1 in 6 cycles  CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery  PD with GEM in 3 cycles. PD with S1 in 3 cycles. PR with HAI (cisplatin). Relapse in 20 months a  PD with weekly paclitaxel in 6 cycles. PR with GEM plus 5-FU in 9 cycles  PR with 5-FU in a year. Sent for conversion surgery. No relapse in 18 months after surgery  SD with GEM in 6months. Excision of new liver metastatic lesion. Relapse in remnant liver in 7 m  PD with S1  N/A
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)  N/A (56)  N/A (57)  10.1111/j.1365-2354.2008.01014. x (58)  10.1186/1477-7819-7-22 (16) 10.1093/jjco/hyp085 (59)  10.1093/jjco/hyp085 (59)	Simon  Yamamoto Nishimizu Morishima Fujii Sorscher  Distler Seki Seki	2011 2012 2011 2010 2010 2009 2009 2009	79 64 64 65 59 48 65 48 67	M M M M M	N/A	Liver metastasis  Peritoneal dissemination  Liver metastasis  Liver metastasis after surgery  Liver metastasis after surgery  Liver metastasis after excision of primary and hepatic lesion  Locally advanced (portal vein)  Liver metastasis after excision of primary and hepatic lesion  Liver metastasis after excision of primary and hepatic lesion  Liver, bone, and lymph-nodes metastasis  Locally advanced (SMA)	S1 GEM plus S1 S1 GEM Paclitaxel 5-FU GEM GEM GEM	CR PR CR PD PD PR SD PD	neurological disorders and relapse. PR with FOLFOX in 12 cycles  CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years  PR with GEM plus S1 in 6 cycles  CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery  PD with GEM in 3 cycles. PD with S1 in 3 cycles. PR with HAI (cisplatin). Relapse in 20 months at PD with weekly paclitaxel in 6 cycles. PR with GEM plus 5-FU in 9 cycles  PR with 5-FU in a year. Sent for conversion surgery. No relapse in 18 months after surgery  SD with GEM in 6months. Excision of new liver metastatic lesion. Relapse in remnant liver in 7 metastatic lesion.
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)  N/A (56)  N/A (57)  10.1111/j.1365-2354.2008.01014.  x (58)  10.1186/1477-7819-7-22 (16)  10.1093/jjco/hyp085 (59)	Simon  Yamamoto Nishimizu Morishima Fujii Sorscher  Distler Seki Seki	2011 2012 2011 2010 2010 2009 2009 2009	79 64 64 65 59 48 65 48	M M M M M	N/A N/A N/A N/A N/A N/A	Liver metastasis  Peritoneal dissemination Liver metastasis Liver metastasis after surgery Liver metastasis after surgery Liver metastasis after excision of primary and hepatic lesion Locally advanced (portal vein) Liver metastasis after excision of primary and hepatic lesion Liver, bone, and lymph-nodes metastasis	S1 GEM plus S1 S1 GEM Paclitaxel 5-FU GEM GEM	CR PR CR PD PD PR SD PD SD	neurological disorders and relapse. PR with FOLFOX in 12 cycles  CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years  PR with GEM plus S1 in 6 cycles  CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery  PD with GEM in 3 cycles. PD with S1 in 3 cycles. PR with HAI (cisplatin). Relapse in 20 months at  PD with weekly paclitaxel in 6 cycles. PR with GEM plus 5-FU in 9 cycles  PR with 5-FU in a year. Sent for conversion surgery. No relapse in 18 months after surgery  SD with GEM in 6months. Excision of new liver metastatic lesion. Relapse in remnant liver in 7 mc  PD with S1  N/A  PD with GEM in 2 cycles. PR with S1 in 4 cycles. PD with additional S1 in 4 cycles  SD with GEM in 6 cycles. New lesion in liver. PD with S1  SD with GEM plus IRI in 7 cycles. SD with GEM plus oxaliplatin in 16 cycles. PD with docetaxel p
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)  N/A (56)  N/A (57)  10.1111/j.1365-2354.2008.01014.  x (58)  10.1186/1477-7819-7-22 (16)  10.1093/jjco/hyp085 (59)  10.1093/jjco/hyp085 (59)  10.1093/jjco/hyp085 (59)  10.1093/jjco/hyp085 (59)  N/A (20)	Simon  Yamamoto Nishimizu Morishima Fujii Sorscher  Distler Seki Seki Seki Seki Antonie	2011 2012 2011 2010 2010 2009 2009 2009	79 64 64 65 59 48 65 48 67 67 61 44	M M M M M M F F	N/A	Liver metastasis  Peritoneal dissemination  Liver metastasis  Liver metastasis after surgery  Liver metastasis after surgery  Liver metastasis after excision of primary and hepatic lesion  Locally advanced (portal vein)  Liver metastasis after excision of primary and hepatic lesion  Liver, bone, and lymph-nodes metastasis  Locally advanced (SMA)  Locally advanced (SMA)  Liver metastasis	S1 GEM plus S1 S1 GEM Paclitaxel 5-FU GEM GEM GEM	CR PR CR PD PD PD SD SD SD	neurological disorders and relapse. PR with FOLFOX in 12 cycles  CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years  PR with GEM plus S1 in 6 cycles  CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery  PD with GEM in 3 cycles. PD with S1 in 3 cycles. PR with HAI (cisplatin). Relapse in 20 months aft  PD with weekly paclitaxel in 6 cycles. PR with GEM plus 5-FU in 9 cycles  PR with 5-FU in a year. Sent for conversion surgery. No relapse in 18 months after surgery  SD with GEM in 6months. Excision of new liver metastatic lesion. Relapse in remnant liver in 7 mc  PD with S1  N/A  PD with GEM in 2 cycles. PR with S1 in 4 cycles. PD with additional S1 in 4 cycles  SD with GEM in 6 cycles. New lesion in liver. PD with S1  SD with GEM plus IRI in 7 cycles. SD with GEM plus oxaliplatin in 16 cycles. PD with docetaxel p capecitabine (6 cycles). PD with FORFIRI. PD with GEM plus erlotinib
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)  N/A (56)  N/A (57)  10.1111/j.1365-2354.2008.01014.  x (58)  10.1186/1477-7819-7-22 (16)  10.1093/jjco/hyp085 (59)  10.1093/jjco/hyp085 (59)  10.1093/jjco/hyp085 (59)  10.1093/jjco/hyp085 (59)  N/A (20)  10.1159/000107397 (19)	Simon  Yamamoto Nishimizu Morishima Fujii Sorscher  Distler Seki Seki Seki Seki Antonie Harten	2011 2012 2011 2010 2010 2009 2009 2009 2009 2009 2009 2007	79 64 64 65 59 48 65 48 67 67 61 44	M M M M M M F F	N/A	Liver metastasis  Peritoneal dissemination Liver metastasis Liver metastasis after surgery Liver metastasis after surgery Liver metastasis after excision of primary and hepatic lesion Locally advanced (portal vein) Liver metastasis after excision of primary and hepatic lesion Liver, bone, and lymph-nodes metastasis Locally advanced (SMA) Locally advanced (SMA) Liver metastasis  Liver metastasis	S1 GEM plus S1 S1 GEM Paclitaxel  5-FU GEM	CR PR CR PD PD PR SD PD SD PR	neurological disorders and relapse. PR with FOLFOX in 12 cycles  CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years  PR with GEM plus S1 in 6 cycles  CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery  PD with GEM in 3 cycles. PD with S1 in 3 cycles. PR with HAI (cisplatin). Relapse in 20 months aff  PD with weekly paclitaxel in 6 cycles. PR with GEM plus 5-FU in 9 cycles  PR with 5-FU in a year. Sent for conversion surgery. No relapse in 18 months after surgery  SD with GEM in 6 months. Excision of new liver metastatic lesion. Relapse in remnant liver in 7 mc  PD with S1  N/A  PD with GEM in 2 cycles. PR with S1 in 4 cycles. PD with additional S1 in 4 cycles  SD with GEM in 6 cycles. New lesion in liver. PD with S1  SD with GEM plus IRI in 7 cycles. SD with GEM plus oxaliplatin in 16 cycles. PD with docetaxel pl  capecitabine (6 cycles). PD with FORFIRI. PD with GEM plus erlotinib  PR with GEM plus mitomycin in 3 months and additional GEM in 3 months
10.2169/internalmedicine.51.6536 (53)  10.3109/0284186X.2011.617388 (54)  N/A (29)  N/A (55)  N/A (56)  N/A (57)  10.1111/j.1365-2354.2008.01014.  x (58)  10.1186/1477-7819-7-22 (16)  10.1093/jjco/hyp085 (59)  10.1093/jjco/hyp085 (59)  10.1093/jjco/hyp085 (59)  10.1093/jjco/hyp085 (59)  N/A (20)	Simon  Yamamoto Nishimizu Morishima Fujii Sorscher  Distler Seki Seki Seki Seki Antonie	2011 2012 2011 2010 2010 2009 2009 2009	79 64 64 65 59 48 65 48 67 67 61 44	M M M M M M F F	N/A	Liver metastasis  Peritoneal dissemination  Liver metastasis  Liver metastasis after surgery  Liver metastasis after surgery  Liver metastasis after excision of primary and hepatic lesion  Locally advanced (portal vein)  Liver metastasis after excision of primary and hepatic lesion  Liver, bone, and lymph-nodes metastasis  Locally advanced (SMA)  Locally advanced (SMA)  Liver metastasis	S1 GEM plus S1 S1 GEM Paclitaxel 5-FU GEM GEM GEM	CR PR CR PD PD PR SD PD SD SD SD PR SD	neurological disorders and relapse. PR with FOLFOX in 12 cycles  CR with S1 in 5 cycles. Sent for conversion surgery. No relapse in 2 years  PR with GEM plus S1 in 6 cycles  CR with S1 in 24 cycles. Excision of multiple right liver tumor. Relapse in 8 months after surgery  PD with GEM in 3 cycles. PD with S1 in 3 cycles. PR with HAI (cisplatin). Relapse in 20 months aft  PD with weekly paclitaxel in 6 cycles. PR with GEM plus 5-FU in 9 cycles  PR with 5-FU in a year. Sent for conversion surgery. No relapse in 18 months after surgery  SD with GEM in 6months. Excision of new liver metastatic lesion. Relapse in remnant liver in 7 mc  PD with S1  N/A  PD with GEM in 2 cycles. PR with S1 in 4 cycles. PD with additional S1 in 4 cycles  SD with GEM in 6 cycles. New lesion in liver. PD with S1  SD with GEM plus IRI in 7 cycles. SD with GEM plus oxaliplatin in 16 cycles. PD with docetaxel plus capecitabine (6 cycles). PD with FORFIRI. PD with GEM plus erlotinib

CA, celiac artery; CR, complete response; F, female; FFX, FORFIRINOX; GEM, gemcitabine; GnP, gemcitabine plus nab-paclitaxel; HAI, hepatic arterial infusion; IRI, irinotecan; M, male; mFFX, modified FORFIRINOX; MSIXEROX, capecitabine and oxaliplatin; N/A, not available; PD, progressive disease; PR, partial response; S1, tegafur/gimeracil/oteracil; SD, stable disease; SMA, superior mesenteric artery; TAE, transcatheter arterial embolization.

## References

- 46. Fontenot J, Spieler B, Hudson C, et al. Corrigendum to "Pancreatic acinar cell carcinoma-literature review and case report of a 56-year-old man presenting with abdominal pain" [Radiol Case Rep 15 (2020) 39-43]. Radiol Case Rep 2019;15:306-7. Erratum for: Radiol Case Rep 2019;15:39-43.
- 47. Yoshida N, Kanno A, Masamune A, et al. Pancreatic Acinar Cell Carcinoma with Multiple Liver Metastases Effectively Treated by S-1 Chemotherapy. Intern Med 2018;57:3529-35.
- 48. Kanematsu K, Suematsu Y, Yamagishi S, et al. Prolonged Survival Achieved with Surgical Resection and Multidisciplinary Therapy for Acinar Cell Carcinoma of the Pancreas with Liver Metastases. Gan To Kagaku Ryoho 2018;45:312-4.
- 49. Luo Y, Hu G, Ma Y, et al. Acinar cell carcinoma of the pancreas presenting as diffuse pancreatic enlargement: Two case reports and literature review. Medicine (Baltimore) 2017;96:e7904.
- Ploquin A, Baldini C, Vuagnat P, et al. Prolonged Survival in a Patient with a Pancreatic Acinar Cell Carcinoma. Case Rep Oncol 2015;8:447-50.
- 51. Callata-Carhuapoma HR, Pato Cour E, Garcia-Paredes B, et al. Pancreatic acinar cell carcinoma with bilateral ovarian metastases, panniculitis and polyarthritis treated with FOLFIRINOX chemotherapy regimen. A case report and review of the literature. Pancreatology 2015;15:440-4.
- 52. Schempf U, Sipos B, König C, et al. FOLFIRINOX as first-line treatment for unresectable acinar cell carcinoma of the pancreas: a case report. Z Gastroenterol 2014;52:200-3.
- 53. Hiraoka A, Nakahara H, Kawasaki H, et al. Huge

- pancreatic acinar cell carcinoma with high levels of AFP and fucosylated AFP (AFP-L3). Intern Med 2012;51:1341-9.
- 54. Simon M, Bioulac-Sage P, Trillaud H, et al. FOLFOX regimen in pancreatic acinar cell carcinoma: case report and review of the literature. Acta Oncol 2012;51:403-5.
- 55. Nishimizu T, Minemura M, Kajiura S, et al. A case of pancreatic acinar cell carcinoma with a giant liver metastasis successfully treated with combination of gemcitabine and peroral S-1. Gan To Kagaku Ryoho 2011;38:309-12.
- Morishima K, Hyodo M, Nihei Y, et al. A case of acinar cell carcinoma of pancreas with liver metastases treated effectively by S-1. Gan To Kagaku Ryoho 2010;37:127-9.
- 57. Fujii M, Sato H, Ogasawara T, et al. A case of liver metastasis of pancreatic acinar cell carcinoma treated with S-1 and intra-arterial CDDP combination therapy. Gan To Kagaku Ryoho 2010;37:1987-90.
- 58. Sorscher SM. Metastatic acinar cell carcinoma of the pancreas responding to gemcitabine, 5-fluorouracil and leucovorin therapy: a case report. Eur J Cancer Care (Engl) 2009;18:318-9.
- 59. Seki Y, Okusaka T, Ikeda M, et al. Four cases of pancreatic acinar cell carcinoma treated with gemcitabine or S-1 as a single agent. Jpn J Clin Oncol 2009;39:751-5.
- 60. Mori S, Kondoh S, Ryozawa S, et al. A case of AFP producing pancreatic acinar cell carcinoma diagnosed by EUS FNA and treated by intraarterial injection chemotherapy. Nihon Shokakibyo Gakkai Zasshi 2004;101:177-82.
- 61. Beltraminelli HS, Buechner SA, Häusermann P. Pancreatic panniculitis in a patient with an acinar cell cystadenocarcinoma of the pancreas. Dermatology 2004;208:265-7.