

Table S1 The baseline characteristics of different groups

Variable	Total sample	≤6 days (N=83)	7–10 days (N=99)	>10 days (N=100)	P value
Underlying disease	7 (2.48%)	4 (4.82%)	3 (3.03%)	0 (0.00%)	0.072
Neutrophil granulocytes [†] (×10 ⁹ /L)	2.66 (1.79–4.23)	2.59 (1.59–3.65)	2.54 (1.47–4.54)	2.85 (2.00–4.31)	0.253
Red blood cells [†] (×10 ¹² /L)	4.67±0.48	4.63±0.48	4.68±0.51	4.71±0.44	0.705
Hemoglobin [†] (g/L)	127.13±11.23	127.65±10.81	127.46±11.36	126.21±11.65	0.540
Platelets [†] (×10 ⁹ /L)	280.42±86.82	269.73±90.79	281.54±79.39	291.12±89.46	0.283
C-reactive protein [†] (mg/dL)	0.52 (0.48–3.83)	0.51 (0.49–3.73)	0.52 (0.47–4.01)	0.51 (0.47–3.08)	0.812
Albumin [†] (g/L)	44.10±2.75	44.21±2.55	43.73±2.84	44.37±2.88	0.309
Alanine aminotransferase [†] (U/L)	33.00 (26.50–41.50)	30.50 (25.00–42.00)	33.00 (27.00–40.50)	35.00 (27.00–40.00)	0.360
Aspartate aminotransferase [†] (U/L)	13.00 (11.00–17.00)	13.00 (11.00–18.00)	13.00 (11.00–17.00)	12.00 (11.00–15.00)	0.654
Creatinine [†] (umol/L)	36.79±10.69	38.21±10.53	36.80±10.20	35.18±11.30	0.097

Data are presented as n (%), median (range), or mean ± SD. [†] Data on the neutrophil granulocytes, red blood cells, hemoglobin, platelets, and C-reactive protein were missing for 52 patients. Data on albumin, alanine aminotransferase, aspartate aminotransferase, and creatinine were missing for 59 patients.