

**Table S1** Primers used for reverse transcription-quantitative PCR

Gene	Forward primer	Reverse primer
<i>β-actin</i>	AGTGTGACGTTGACATCCGT	GCAGCTCAGTAACAGTCCGC
<i>Il6</i>	TAGTCCTTCCTACCCCAATTTCC	TAGTCCTTCCTACCCCAATTTCC
<i>Tnfα</i>	AAAATTCGAGTGACAAGCCTGTAG	CCCTTGAAGAGAACCTGGGAGTAG
<i>Il1b</i>	GACCTTCCAGGATGAGGACA	AGCTCATATGGGTCCGACAG
<i>iNOS</i>	ACATCGACCCGTCCACAGTAT	CAGAGGGGTAGGCTTGTCTC
<i>Arg1</i>	CTCCAAGCCAAAGTCCTTAGAG	AGGAGCTGTCATTAGGGACATC
<i>Ccl2</i>	CCTGCTGCTACTCATTAC CA	ATTCCTTCTTGGGGTCAG CA

**Table S2** Baseline characteristics of collected subjects

Characteristics	Sepsis patients (n=55)	Non-septic patients (n=18)	P
Age (years), mean ± SD	68.24±16.43	60.56±14.47	0.08
BMI (kg/m <sup>2</sup> ), mean ± SD	21.65±2.28	22.61±2.65	0.13
Sex, n (%)			0.84
Male	43 (78.2)	13 (72.2)	
Female	12 (21.8)	5 (27.8)	
The origin of sepsis, n (%)			
Lung infection	8 (14.5)		
Abdominal infection	45 (81.8)		
Intracranial infection	2 (3.6)		