

Supplementary

Table S1 Patient and tumor characteristics for the external validation cohort

| Characteristics | All external validation cohort | | |
|-------------------------------------|--------------------------------|--------------------------------|---------------------------------|
| | Sum (n=261) | Medical center 1 (n=94), n (%) | Medical center 2 (n=167), n (%) |
| Age (years), mean ± SD | 53.16±8.07 | 53.05±9.13 | 53.36±8.46 |
| Tumor size (cm), mean ± SD | 3.55±1.86 | 3.86±2.34 | 3.24±1.57 |
| Tumor histological subtype | | | |
| I | 220 | 80 (85.11) | 140 (83.83) |
| II | 15 | 6 (6.38) | 9 (5.39) |
| Mixed (I + II) | 26 | 8 (8.51) | 18 (10.78) |
| Tumor grade (differentiation) | | | |
| 1–2 (well/moderate differentiation) | 176 | 63 (67.02) | 113 (67.66) |
| 3 (poor differentiation) | 85 | 31 (32.98) | 54 (32.34) |
| Depth of tumor invasion | | | |
| <1/2 | 134 | 44 (46.81) | 90 (53.89) |
| ≥1/2 | 127 | 50 (53.19) | 77 (46.11) |
| LVSI | | | |
| Yes | 119 | 48 (51.06) | 71 (42.51) |
| No | 142 | 46 (48.94) | 96 (57.49) |
| Lymph node metastasis | | | |
| Yes | 114 | 41 (43.62) | 73 (43.71) |
| No | 147 | 53 (56.38) | 94 (56.29) |
| FIGO stage | | | |
| I | 130 | 41 (43.62) | 89 (53.29) |
| II | 46 | 7 (7.45) | 39 (23.35) |
| III | 70 | 37 (39.36) | 33 (19.76) |
| IV | 15 | 9 (9.57) | 6 (3.59) |

SD, standard deviation; LVSI, lymph vascular space invasion; FIGO, International Federation of Gynecology and Obstetrics.