

Table S1 Association of PCAT_{MA} with different demographic factors in subgroups stratified by sex, BMI and age

Characteristics	PCAT _{MA}					
	Sex		BMI (kg/m ²)		Age (years)	
	Women	Men	≥24.91	<24.91	≥55	<55
Age (years)						
rs	0.013	-0.007	-0.020	0.021	-	-
P value	0.790	0.900	0.695	0.679	-	-
Gender						
rs	-	-	0.019	0.021	0.009	0.032
P value	-	-	0.707	0.676	0.863	0.529
BMI (kg/m ²)						
rs	0.053	0.051	-	-	0.031	0.074
P value	0.271	0.330	-	-	0.525	0.144
Smoking						
rs	0.101*	-0.046	0.016	0.036	0.015	0.046
P value	0.036	0.370	0.751	0.467	0.762	0.358
Hypertension						
rs	0.048	0.092	0.144**	-0.009	0.064	0.076
P value	0.315	0.077	0.004	0.851	0.193	0.130
Hyperglycaemia						
rs	0.008	-0.031	0.036	-0.053	-0.011	-0.006
P value	0.873	0.549	0.469	0.286	0.832	0.900
Dyslipidemia						
rs	0.328**	0.339**	0.342**	0.326**	0.367**	0.298**
P value	<0.001	<0.001	<0.001	<0.001	<0.001	<0.001

** , significant association at the 0.01 level (two-tailed); * , significant association at the 0.05 level (two-tailed). PCAT_{MA}, mean pericoronary adipose tissue attenuation; BMI, body mass index.