

Table S1 Absolute lymphocyte counts during treatment

Parameter	Non-T α 1 group (N=48)	Short-term T α 1 group (N=101)	Long-term T α 1 group (N=47)	P value
Mid-CCRT ALC				
G0-2	41 (85.4%)	84 (84.8%)	38 (80.9%)	0.79
G3-4	7 (14.6%)	15 (15.2%)	9 (19.1%)	
2 m post-CCRT ALC				
G0	14 (34.1%)	39 (41.5%)	23 (51.1%)	0.28
G1-3	27 (65.9%)	55 (58.5%)	22 (48.9%)	
6 m post-CCRT ALC				
G0	14 (45.2%)	47 (69.1%)	31 (77.5%)	0.01
G1-3	17 (55.8%)	21 (30.9%)	9 (22.5%)	
9 m post-CCRT ALC				
G0	10 (55.6%)	39 (75.0%)	25 (83.3%)	0.10
G1-3	8 (44.4%)	13 (25.0%)	5 (16.7%)	
12 m post-CCRT ALC				
G0	11 (61.1%)	32 (78.0%)	21 (80.8%)	0.28
G1-3	7 (38.9%)	9 (22.0%)	5 (19.2%)	
15 m post-CCRT ALC				
G0	4 (57.1%)	25 (86.2%)	21 (80.8%)	0.22
G1-3	3 (42.9%)	4 (13.8%)	5 (19.2%)	
ALC at nadir (k/ μ L)				
Median (range)	0.74 (0.36-1.41)	0.70 (0.15-1.95)	0.73 (0.16-1.54)	0.89