

Table S1 Plasma albumin decline between the 2 study groups

Characteristics	Clamping group			Control group			P ¹	P ²	P ³
	POD 1 (n=85)	Discharge (n=85)	1 mo (n=61)	POD1 (n=82)	Discharge (n=82)	1 mo (n=59)			
Albumin declination, mean ± SD, g/L	7.3±3.1	7.7±2.9	1.0±3.2	7.4±3.0	9.0±5.2	1.6±2.7	0.849	0.040	0.212

P¹, P value between 2 group at POD1; P², P value between 2 group at discharge; P³, P value between 2 group at 1 mo. POD, postoperative day; mo, month; SD, standard deviation.

Table S2 Hemoglobin decline between the 2 study groups

Characteristics	Clamping group			Control group			P ¹	P ²	P ³
	POD 1 (n=85)	Discharge (n=85)	1 mo (n=58)	POD1 (n=83)	Discharge (n=82)	1 mo (n=60)			
Hemoglobin declination, mean ± SD, g/L	10.3±8.3	16.8±9.0	2.7±8.4	10.0±7.6	20.4±14.4	2.5±6.8	0.856	0.050	0.883

P¹, P value between 2 group at POD1; P², P value between 2 group at discharge; P³, P value between 2 group at 1 mo. POD, postoperative day; mo, month; SD, standard deviation.