

Supplementary

Table S1 Scoring criteria for the respiratory symptom subscales

Item	No (score 0)	Mild (score 1)	Moderate (score 2)	Severe (score 3)
Cough	None	Intermittent cough, not affecting normal work or study	Between mild and severe	Frequent day and night coughing, or bouts of coughing that interfered with work and sleep
Sputum amount	None	Cough with sputum, occasional sputum in both lungs	Between mild and severe	Cough with sputum, both lungs full of sputum
Expectoration difficulty	None	Easy expectoration	Somewhat difficult expectoration	Difficult expectoration
Wheezing rate	None	Occasional wheezing rate	Scattered wheezing rate	Consistent wheezing rate




Figure S1 Scoring criteria of the VAS for self-reported symptom severity. VAS, visual analog scale.

Table S2 Comparison of vital signs at D7 between the AHCHOS and AHI groups.

Items	Total (n=407)	AHI (n=153)	AHCHOS (n=254)	P value
Heart rate				0.63
n (missing)	375 (32.0)	137 (16.0)	238 (16.0)	
Median (IQR)	116.00 (108.00–124.00)	119.00 (108.00–125.00)	116.00 (107.00–124.00)	
Systolic blood pressure				0.54
n (missing)	54 (353.0)	28 (125.0)	26 (228.0)	
Median (IQR)	90.00 (85.00–99.00)	92.00 (85.00–99.50)	90.00 (85.00–99.00)	
Diastolic blood pressure				0.71
n (missing)	54 (353.0)	28 (125.0)	26 (228.0)	
Median (IQR)	58.00 (52.00–60.00)	59.00 (50.00–61.00)	58.00 (56.00–60.00)	

Data are presented as n (%) unless otherwise indicated. AHI, ambroxol hydrochloride injection; AHCHOS, ambroxol hydrochloride and clenbuterol hydrochloride oral solution; IQR, interquartile range.