

Table S1 Blood results of laboratory tests in 12 children with EIMS

Case	C-reactive protein (mg/L)	White blood cell ($10^9/L$)	Red blood cell ($10^{12}/L$)	Hemoglobin (g/L)	Blood platelet ($10^9/L$)	Plateletcrit (%)	Neutrophil count ($10^9/L$)	Lymphocyte count ($10^9/L$)	Prealbumin (mg/L)	Albumin (g/L)	Albumin-globulin ratio
1	26	8.13	4.45	119	276	0.48	3.22	1.67	70	32.8	1.18
2	90	9.02	3.79	90	904	0.80	3.10	0.43	167	38.3	1.50
3	47	12.75	5.50	131	380	0.40	8.21	0.34	165	31.3	2.10
4	20	5.87	4.64	109	466	0.39	2.94	2.23	91	32.8	0.97
5	<8	6.35	3.61	73	1,094	0.87	4.03	1.87	65	29.8	0.53
6	53	12.99	4.58	125	413	0.40	10.2	1.93	70	31.3	1.09
7	<8	8.96	4.75	141	243	0.24	6.38	29.7	192	46.4	1.42
8	<8	11.37	4.71	130	404	0.40	5.19	3.81	NA	40.0	1.74
9	33	9.47	4.08	102	610	0.54	5.21	2.64	53	34.5	1.14
10	10	12.46	4.84	133	579	0.63	7.35	3.18	68	39.4	1.41
11	78.66	28.98	3.34	78	653	0.68	25.18	2.00	50	31.9	0.94
12	105	13.48	3.52	73	1,061	0.96	8.22	2.70	NA	28.8	0.88

C-reactive protein normal value: <8 mg/L; white blood cell normal value: $(4.0-10.0) \times 10^9/L$; red blood cell normal value: $(4.0-5.0) \times 10^{12}/L$; hemoglobin normal value: 118–156 g/L; blood platelet normal value: $(190-524) \times 10^9/L$; plateletcrit normal value: 0.19–0.34%; neutrophil count normal value: $(0.8-5.8) \times 10^9/L$; lymphocyte count normal values: $(2.4-8.7) \times 10^9/L$; prealbumin normal value: 150–400 mg/L; albumin normal value: 39–54 g/L; albumin-globulin ratio normal value: 1.4–3.9. EIMS, epithelioid inflammatory myofibroblast sarcoma; NA, data not available.