Table S1 Doses of drugs for Beijing Children's Hospital (BCH)-rhabdomyosarcoma (RMS)-2006

| Drugs | Doses |
|-----------------------|---|
| Vincristine (V) | >3 years, 1.5 mg/m ² |
| | 1–3 years, 0.05 mg/kg |
| | <1 year, 0.025 mg/kg |
| | Top dose (2 mg) used on Days 1, 8 and 15 |
| Dactinomycin (A) | ≥1 year, 0.045 mg/kg |
| | <1 year, 0.025 mg/kg |
| | On Day 1 (maximum dose, 2.5 mg) |
| Topotecan (T)* | 1.5 mg/m ² on Days 1–3 |
| Cyclophosphamide (C)* | 2.2 g/m ² on Day 1 for the vincristine, dactinomycin, and cyclophosphamide (VAC) protocol |
| | 750 mg/m ² /dose on Days 1–2 for the vincristine, topotecan, and cyclophosphamide (VTC) protocol |

*, for children aged under 1 year or weighing less than 10 kilograms, drugs were converted into body weight for calculation.

| Drugs | Doses |
|-------------------------------|---|
| Vincristine (V) | The same as in the Beijing Children's Hospital (BCH)-RMS-2006 |
| Dactinomycin (A) | The same as in the BCH-RMS-2006 |
| Dactinomycin (A) [#] | 1.5 mg/m ² on Day 1 |
| Cyclophosphamide (C) | ≥1 year, 1.2 g/m ² |
| | <1 year, 40 mg/kg |
| | On Day 1 (with mesna uroprotection) |
| Irinotecan (Irin) | 50 mg/m²/dose, on Days 1–5, with a maximum of 100 mg/dose |
| Doxorubicin (D) | ≥1 year, 30 mg/m² |
| | <1 year, 1 mg/kg |
| | On Days 1–2 |
| Doxorubicin (D) [#] | 25 mg/m ² |
| | On Days 1–2 |
| Ifosfamide (I) | ≥1 year, 1800 mg/m² |
| | <1 year,60 mg/kg |
| | On Days 1–5 (with mesna uroprotection) |
| lfosfamide (I) [#] | 3 g/m ² , on Days 1–5 (with mesna uroprotection) |
| Etoposide (E) | ≥1 year, 100 mg/m² |
| | <1 year, 5 mg/kg |
| | On Days 1–5 |
| Etoposide (E) [#] | 150 mg/m ² |
| | On Days 1–3 |
| Carboplatin [#] | 560 mg/m ² |
| | On Day 1 |

*, drug doses in the Central Nervous Invasion (CNI) group in the CCCG-RMS-2016 protocol.