



# H3K27ac-activated lncRNA *KTN1-AS1* aggravates tumor progression by miR-505-3p/*ZNF326* axis in ovarian cancer

Xiaohui Xie<sup>1#</sup>, Qirong Wen<sup>1#</sup>, Xu Yang<sup>2</sup>, Weiwei Chen<sup>1</sup>, Yaqiong Liu<sup>1</sup>, Wenjie Liu<sup>1</sup>, Tiansong Zhang<sup>1</sup>, Cuiyi Xu<sup>3</sup>, Kun Shi<sup>1</sup>

<sup>1</sup>Department of Gynecology and Obstetrics, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, China; <sup>2</sup>Department of Obstetrics and Gynecology, The Fifth Affiliated People's Hospital of Chengdu University of Traditional Chinese Medicine, Chengdu, China; <sup>3</sup>Department of Operation Room Nursing, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, China

**Contributions:** (I) Conception and design: K Shi; (II) Administrative support: K Shi; (III) Provision of study materials or patients: X Xie, Q Wen, X Yang; (IV) Collection and assembly of data: X Xie, Q Wen; (V) Data analysis and interpretation: Q Wen; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

<sup>#</sup>These authors contributed equally to this work.

**Correspondence to:** Kun Shi. Department of Gynecology and Obstetrics, Guangzhou Women and Children's Medical Center, Guangzhou Medical University, Guangzhou, China. Email: shikun28@hotmail.com.

**Background:** Numerous studies indicate that long noncoding RNA (lncRNA) is aberrantly expressed in ovarian cancer (OC). Our research investigated the regulatory role of lncRNA *KTN1* antisense RNA1 (*KTN1-AS1*) in the progression of OC through the miR-505-3p/*ZNF326* axis.

**Methods:** Expression of *KTN1-AS1*, microRNA-505-3p (*miR-505-3p*), and zinc-finger protein-326 (*ZNF326*) in OC was evaluated by using RT-qPCR analysis. The biological function of *KTN1-AS1* was inspected using the loss-of-function assay. Luciferase reporter assay and RIP assay were performed to determine the competitive endogenous RNA (ceRNA) network of *KTN1-AS1/miR-505-3p/ZNF326*.

**Results:** The data showed that *KTN1-AS1* and *ZNF326* had a high expression in OC than in the normal tissue, and *miR-505-3p* exhibited a low expression in OC than in the normal tissue. The knockdown of *KTN1-AS1* caused an inhibition in OC cell proliferation, migration, and invasion, and promoted cell apoptosis. In terms of mechanical exploration, *KTN1-AS1* was transcriptionally activated by histone H3 on lysine 27 acetylation (H3K27ac) at the promoter region, and *KTN1-AS1* increased *ZNF326* expression by competitively adsorbing *miR-505-3p*.

**Conclusions:** This study indicated that H3K27ac-induced lncRNA *KTN1-AS1* expression, and facilitated proliferation, migration, and invasion of OC cells by the *KTN1-AS1/miR-505-3p/ZNF326* axis.

**Keywords:** H3K27ac; *KTN1-AS1*; *miR-505-3p*; *ZNF326*; ovarian cancer (OC)

Submitted Nov 24, 2021. Accepted for publication May 20, 2022. This article was updated on July 01, 2025.

The original version is available at: <http://dx.doi.org/10.21037/atm-22-443>

doi: 10.21037/atm-22-443

## Introduction

Ovarian cancer (OC) is a common gynecologic cancer, and its occurrence rate is the seventh highest of all malignancies in females (1). Importantly, OC is also the fifth leading reason for tumor-relevant deaths in females (2). Reformative techniques in diagnosis and therapy have

decreased the incidence and mortality rate of patients with OC. Nevertheless, the rate of 5-year survival rates remains unsatisfying in advanced OC patients because of extensive metastases, drug resistance, and relapse of OC (3). Therefore, it is essential to perform further researches for the regulatory mechanism of OC progression and development.

Long noncoding RNA (lncRNA) is a cluster of transcripts

( $\geq 200$  nucleotides) with little or without capacity in coding proteins (4). Previously, these transcripts were considered as transcriptional *noise*; however, emerging research found that some of these transcripts exhibited essential regulatory properties in various physiological or pathological processes (5,6). Based on some studies, lncRNAs were widely reported to modulate genomic imprinting (7), X chromosome inactivation (8), stem cell differentiation (9), and cancer metastasis (10-12). The modulatory roles of lncRNAs have been reported in colon cancer (13), gastric cancer (14), and cervical cancer (15). Recent research also found that a series of lncRNAs with an abnormal expression exerted important functions in OC, such as *PTAR* (16), *PVT1* (17), *MALAT1* (18), and *HIST2* (19). Nonetheless, effects of most lncRNAs remain largely unclear in OC, and there is need to study these effects further.

*KTN1* antisense RNA 1 (*KTN1-AS1*) is a newly recognized tumor-associated lncRNA that is known to play a regulatory role in diverse cell types. In hepatocellular carcinoma (HCC), *KTN1-AS1* sponges *miR-23c* to affect *ERBB2IP* expression, thereby promoting the growth of HCC (20). *KTN1-AS1* also expresses at a high level and is associated with unsatisfying clinical outcomes in lung cancer patients (21). In addition, *KTN1-AS1* has shown an upregulated expression level and facilitates the migration and invasion of bladder cancer cells (22). However, the expression and biological function of *KTN1-AS1* were not inspected in relation to OC.

H3K27ac is associated with gene activation and is mainly concentrated in the enhancer and promoter regions. When the promoter or enhancer region is modified with H3K27ac, the promoter or enhancer will be activated to promote gene expression.

In our study, we explored the function and molecular mechanism of *KTN1-AS1* in OC cells and found the *KTN1-AS1/miR-505-3p/ZNF326* axis. This finding offers a promising insight into the treatment of OC. We present the following article in accordance with the MDAR reporting checklist (available at <https://atm.amegroups.com/article/view/10.21037/atm-22-443/rc>).

## Methods

### *Clinical specimens*

Sixty-four OC and ten normal ovarian tissues were collected from patients at Guangzhou Women and

Children's Medical Center, Guangzhou Medical University. An informed consent form was signed by each patient, and none of the patients accepted anti-cancer therapy before surgery. All collected tissues were preserved at  $-80^{\circ}\text{C}$  in liquid nitrogen. The study was approved by the Ethics Committee of Guangzhou Women and Children's Medical Center, Guangzhou Medical University (approval No. 2020-KY-021-03). The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013).

### *Cell lines and reagent*

IOSE-80 (the human ovarian cell line) and A2780, OVCAR-3, SKOV3 and TOV112D (OC cell lines) were provided by ATCC (Manassas, VA, USA). Dulbecco's modified Eagle's medium (DMEM; HyClone, Logan, UT, USA), which contains 10% fetal bovine serum (FBS; HyClone, Logan, UT, USA), was utilized to culture these cell lines at  $37^{\circ}\text{C}$  with 5%  $\text{CO}_2$ . The acetylation inhibitor (C646) was commercially obtained from Sigma Chemical (St. Louis, MO, USA).

### *Cell transfection*

The short hairpin RNA (shRNA) against *KTN1-AS1* (*sh-KTN1-AS1#1/2*) or *CBP* (*sh-CBP#1/2*), *miR-505-3p* mimics, and their relative reference (sh-NC and NC mimics) were synthesized by GenePharma (Shanghai, China). To overexpress *ZNF326*, *pcDNA3.1/ZNF326* (GenePharma, Shanghai, China) was synthesized, and the empty *pcDNA3.1* vector was taken as an internal reference. Cell transfection was performed through Lipofectamine 2000 (Invitrogen, Carlsbad, CA, USA). Forty-eight hours later, transfection efficiency was proved by RT-qPCR.

### *RT-qPCR*

TRIzol reagent (Invitrogen, Carlsbad, CA, USA) was used to isolate the total RNA from OC tissues and cells. Then, PrimeScript RT reagent Kit (Takara, Tokyo, Japan) was employed to reverse-transcribe the isolated RNA into complementary DNA (cDNA). On an ABI7300 real-time PCR machine (Applied Biosystems, Carlsbad, CA, USA), SYBR Green PCR Master Mix (Roche, Basel, Switzerland) was used for RT-qPCR to determine relative RNA expression. The appointed genes were normalized to Glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*)

**Table 1** Sequences of the primers and target shRNA

Name	Sequences of the primers
<i>KTN1-AS1-F</i>	GCGAAGCCGTTAGTCCCTTA
<i>KTN1-AS1-R</i>	TTGGGTGAAAGTGGACCTGG
<i>ZNF326-F</i>	AATGAACCCGAACAAAGCCG
<i>ZNF326-R</i>	ACCTCCGAAAGAGTCAAGGC
<i>hsa-miR-505-3p-F</i>	CGTCAACACTTGCTGGTTTCT
<i>CYTH1-F</i>	CACCATGGAGGAGGACGAC
<i>CYTH1-R</i>	CCGTGGGATGTTCTCCAGTT
<i>DNAJB9-F</i>	AGAGCGCCAAATCAAGAAGG
<i>DNAJB9-R</i>	TTCAGCATCCGGGCTCTTATT
<i>U6-F</i>	ATGGACTATCATATGCTTACCGTA
<i>NCK2-F</i>	CACGGCGAGATTTTCATGTGTT
<i>NCK2-R</i>	GCAGGGATTCTCCAAATCCTCA
<i>KTN1-AS1 ShRNA</i>	GACTGTGGATAGAGATAGATAGATTCAAGA GAGACTGTGGATAGAGATAGATTTTTT

(control for lncRNA and mRNA) or *U6* (miRNA control) using the  $2^{-\Delta\Delta C_t}$  method. The primer sequences used in this study are listed in *Table 1*.

### Colony formation assay

Transfected A2780 and OVCAR-3 cells were planted in 6-well plates including a complete medium with 10% fetal bovine serum (FBS). Two weeks later, phosphate-buffered saline (PBS; Thermo Fisher Scientific, Carlsbad, CA, USA) was used to wash the colonies twice. Then, the colonies were fixed by 4% polyoxymethylene and colored with 1% crystal violet. Colonies (exceeding 50 cells) were manually counted.

### Flow cytometry analysis

In 6-well plates, transfected A2780 and OVCAR-3 cells were rinsed in PBS. After trypsinization, the cells were re-suspended in a binding buffer (100  $\mu$ L). FITC-conjugated with Annexin V and PI (1  $\mu$ L; Invitrogen, Carlsbad, CA, USA) was added into the binding buffer. Fifteen minutes later, the apoptotic OC cells were analyzed using flow cytometry (BD Biosciences, NY, USA).

### Wound healing assay

The transfected A2780 and OVCAR-3 cells were seeded in 6-well plates ( $5 \times 10^5$  cells/well). At 80% confluence, a 200  $\mu$ L pipette tip was used in the monolayer of cells to generate wounds. Then, cells were washed with PBS, and the function of genes on cell migration was assessed after adding the medium containing mitomycin C (20  $\mu$ g/mL) without FBS. Photographs were taken to evaluate healing 0 and 24 hours after wounding.

### Transwell assay

Invasion of transfected A2780 and OVCAR-3 cells was assessed by applying a Transwell chamber with Matrigel (Corning, NY, USA). Add a serum-free medium containing  $1 \times 10^5$  cells to the upper chamber, and add a medium containing 10% FBS to the lower chamber, and the medium containing 10% FBS was placed into the lower chamber. This was incubated for 48 hours, then invaded cells were fixed, stained with 1% crystal violet staining solution, then rinsed with PBS and counted with five fields in random.

### ChIP assay

The ChIP Assay Kit (Thermo Fisher Scientific, Carlsbad, CA, USA) was used to examine the ChIP assay in A2780 and OVCAR-3 cells. DNA-protein cross-links were formed after the incubation of A2780 and OVCAR-3 cells with formaldehyde for 10 minutes. Next, cross-linked chromatin DNAs were broken into segments sized 200–1,000 bp using an ultrasound machine. Then, the lysate was immunoprecipitated with anti-H3K27ac (Abcam, Boston, MA, USA), anti-CBP (Abcam, Boston, MA, USA), or IgG antibody (control, Abcam, Boston, MA, USA), and precipitates were recovered for RT-qPCR analysis.

### RNA pull down assay

The protein was extracted from A2780 and OVCAR-3 cells and later treated with purified biotinylated RNA (*KTN1-AS1* biotin probe) and biotinylated NC (*KTN1-AS1* no-biotin probe). The purified biotinylated transcripts were commercially obtained from Sangon Biotech (Shanghai, China) and were used to incubate with cell lysates at 25 °C for 1 hour. Later, streptavidin agarose beads (Invitrogen,

Carlsbad, CA, USA) were applied to isolate biotin-coupled RNA complexes, and RT-qPCR was utilized to analyze miRNAs enrichment in the pull-down products.

### *Luciferase reporter assay*

*KTN1-AS1* full-length and *ZNF326* 3'-UTR of the wild-type (WT) and mutant-type (Mut) *miR-505-3p* binding sites were generated in Genechem Co., Ltd. (Shanghai, China). *KTN1-AS1*-WT/Mut or *ZNF326*-WT/Mut was transfected into A2780 and OVCAR-3 cells with *miR-505-3p* mimics/NC mimics. Luciferase activity was evaluated using Dual-Luciferase Reporter Assay (Promega, Madison, USA) after 48 hours of transfection.

### *RIP assay*

A2780 and OVCAR-3 cells were lysed in the lysis buffer, and the supernatant was collected after the lysate was centrifuged. Later, cell lysate was adopted to incubate with an anti-Ago2 or anti-IgG antibody (Millipore, MO, USA), and then magnetic beads were added for the immunoprecipitate RNA. After purification, RNA was detected by RT-qPCR.

### *Statistical analysis*

Statistical analyses were performed with GraphPad Prism Software (San Diego, CA, USA), and results were shown as mean  $\pm$  standard deviation (SD) based on no less than three repeats. A Student's *t*-test or one-way Analysis of Variance (ANOVA) was used to estimate group difference, and statistical significance was set as  $P < 0.05$ .

## **Results**


### *KTN1-AS1 was upregulated and enhanced cell proliferation and invasion in OC*


To determine the relationship between *KTN1-AS1* and the biological processes of OC, we first detected *KTN1-AS1* expression using RT-qPCR analysis. The results confirmed the increased *KTN1-AS1* expression in OC tissues compared to the expression in adjacent normal tissues (*Figure 1A*). Consistent with this, we compared these results with the IOSE-80 cell line; *KTN1-AS1* was present in a high level in OC cell lines (*Figure 1B*). To examine whether upregulated *KTN1-AS1* impacted OC cell growth, we performed a

loss-of-function assay by constructing shRNA-mediated silencing of *KTN1-AS1* in A2780 and OVCAR-3 cells. The results showed that sh-*KTN1-AS1*#1/2 transfection caused a decrease in *KTN1-AS1* expression (*Figure 1C*). The function of silenced *KTN1-AS1* in cell proliferation was estimated using a colony formation assay, and data showed that *KTN1-AS1* deficiency slowed the proliferation of A2780 and OVCAR-3 cells (*Figure 1D*). Flow cytometry analysis showed that the apoptosis increased in *KTN1-AS1*-silenced A2780 and OVCAR-3 cells (*Figure 1E*). Subsequently, the wound healing assay demonstrated inhibitive migration in the sh-*KTN1-AS1*#1/2 group (*Figure 1F*). Furthermore, OC cell invasion was also suppressed by the deficiency of *KTN1-AS1* in the Transwell assay (*Figure 1G*). Taken together, *KTN1-AS1* was highly expressed and played a facilitative role in OC cell proliferation and invasion.


### *KTN1-AS1 was transcriptionally upregulated by CBP-mediated H3K27ac*

Then, we probed the mechanism of *KTN1-AS1* upregulation in OC. Previous research has shown that histone H3 lysine 27 acetylation (*H3K27ac*) could activate lncRNAs at the transcriptional level (23,24). By using UCSC (<http://genome.ucsc.edu/>), we also found that the *KTN1-AS1* promoter region presented high H3K27ac enrichment (*Figure 2A*). Therefore, we hypothesized that high expression of *KTN1-AS1* was triggered by H3K27ac at its promoter region. To verify this, ChIP assay was carried out to evaluate the H3K27ac level at the *KTN1-AS1* promoter region in OC cells (A2780 and OVCAR-3) and in the IOSE-80 cell line. Results showed that the H3K27ac level in the *KTN1-AS1* promoter region was higher in A2780 and OVCAR-3 cells compared to the IOSE-80 cell line (*Figure 2B*). Then, the histone acetyltransferase (HAT) inhibitor (C646) was used, and we found that *KTN1-AS1* expression was reduced by C646 treatment compared to dimethyl sulfoxide (DMSO) treatment (*Figure 2C*). These results indicated that *KTN1-AS1* was induced by H3K27ac. Considering that *CBP* acts as a key modulator in gene transcription and histone acetylation (25), we examined whether *CBP* was in charge of H3K27ac in the *KTN1-AS1* promoter. Using ChIP analysis, the enrichment of the *KTN1-AS1* promoter in *CBP* precipitates was validated (*Figure 2D*), highlighting the binding of *CBP* to the *KTN1-AS1* promoter. Later, *CBP* expression was knocked down in A2780 and OVCAR-3 cells (*Figure 2E*). We further found that *KTN1-AS1* expression decreased upon *CBP* knockdown





**Figure 1** *KTN1-AS1* exhibited high level and oncogenic property in OC cells. (A) RT-qPCR analysis of *KTN1-AS1* expression in 64 pairs of OC and adjacent normal tissues. (B) RT-qPCR analysis of *KTN1-AS1* expression in OC cell lines (A2780, OVCAR-3, SKOV3, and TOV112D) and normal human ovarian cell line (IOSE-80). (C) *KTN1-AS1* expression was detected in A2780 and OVCAR-3 cells transfected with *KTN1-AS1* shRNAs by qRT-PCR. (D) Colony formation assay in *KTN1-AS1*-SH#1/2 transfected A2780 and OVCAR-3 cells. (E) Flow cytometry analysis used to evaluate the effect of *KTN1-AS1* silencing on cell apoptosis. (F) Wound healing assay used to estimate the migration ability upon *KTN1-AS1* knockdown. Only use 1% crystal violet staining solution for staining. The magnification is 100 $\times$ . (G) Transwell assay used to assess the function of *KTN1-AS1* deficiency in cell invasion. Only use 1% crystal violet staining solution for staining. The magnification is 400 $\times$ . \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ . NC, negative control; OC, ovarian cancer; RT-qPCR, real-time quantitative polymerase chain reaction.




**Figure 2** Transcriptional activation of *KTN1-AS1* by *CBP* (*CREB binding protein*)-mediated H3K27ac (histone H3 on lysine 27 acetylation). (A) UCSC (University of California, Santa Cruz) was used to predict H3K27ac enrichment at *KTN1-AS1* promoter. (B) H3K27ac level on *KTN1-AS1* promoter was determined by ChIP assay in A2780, OVCAR-3 and IOSE-80 cells. A2780 cell, human ovarian cancer epithelial cell line, OVCAR-3 cells, human ovarian adenocarcinoma epithelial cell line, IOSE-80 cells, human normal ovarian epithelial cell line. (C) RT-qPCR result of *KTN1-AS1* expression in C646- or DMSO-treated A2780 and OVCAR-3 cells. C646, a histone acetyltransferase inhibitor. (D) The binding of *CBP* to *KTN1-AS1* promoter was confirmed by ChIP assay. (E) RT-qPCR validated the silence of *CBP* in OC cells. (F) *KTN1-AS1* expression in OC cells upon *CBP* deficiency was detected by RT-qPCR. (G) The function of silenced *CBP* in H3K27ac level on *KTN1-AS1* promoter was determined by ChIP assay. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ . NC, negative control; OC, ovarian cancer; RT-qPCR, real-time quantitative polymerase chain reaction.

in OC cells (Figure 2F). Importantly, we confirmed that *CBP* silencing decreased the H3K27ac level in the *KTN1-AS1* promoter region (Figure 2G). Conclusively, *KTN1-AS1* was induced by *CBP*-mediated H3K27ac in OC.

#### *KTN1-AS1* served as competitive endogenous RNA (ceRNA) for *miR-505-3p*

Later, we investigated the downstream regulatory pattern of *KTN1-AS1* in OC. Previous studies showed that *KTN1-*

*AS1* was involved in the ceRNA network in some cancers (20,21). Thus, we used starBase (<http://starbase.sysu.edu.cn/>) to seek microRNAs (miRNAs) that could theoretically bind to *KTN1-AS1*, and five potential miRNAs were found (Figure 3A). The RNA pull down assay with biotin-labeled *KTN1-AS1* implied that *miR-505-3p* was pulled down by the *KTN1-AS1* biotin probe (Figure 3B). Thus, we selected *miR-505-3p* to conduct subsequent research. *miR-505-3p* expression was examined in OC tissues and cells, and RT-qPCR demonstrated that *miR-505-3p* expressed at a low



**Figure 3** *KTN1-AS1* sponged *miR-505-3p* in OC. (A) The potential miRNAs for *KTN1-AS1* in starBase. (B) The relative enrichment of predicted miRNAs interacting with *KTN1-AS1* was quantified by RT-qPCR after biotinylated-*KTN1-AS1* pull-down experiment in OC cells. (C) *miR-505-3p* expression in OC tissues and cells was examined by RT-qPCR. (D) Overexpression of *miR-505-3p* in A2780 and OVCAR-3 cells was verified by RT-qPCR. (E) *miR-505-3p* binding site and mutated site on *KTN1-AS1*. (F) The luciferase activity of *KTN1-AS1*-WT/Mut in OC cells with *miR-505-3p* mimics was validated by luciferase reporter assay. \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ . NC, negative control; OC, ovarian cancer; RT-qPCR, real-time quantitative polymerase chain reaction.


level (Figure 3C). Furthermore, we elevated *miR-505-3p* expression by transfecting *miR-505-3p* mimics (Figure 3D). The binding sequences of *miR-505-3p* in *KTN1-AS1*-WT and *KTN1-AS1*-Mut are represented in Figure 3E. In addition, we observed that the luciferase activity of the *KTN1-AS1*-WT reporter was reduced, and that of the *KTN1-AS1*-Mut reporter was not affected under the transfection of *miR-505-3p* mimics (Figure 3F). All data showed that *KTN1-AS1* interacted with *miR-505-3p* in OC.


### *ZNF326* was targeted by *miR-505-3p*

Subsequently, the target gene of *miR-505-3p* was inspected

to further support the ceRNA hypothesis. By using starBase (CLIP Data  $\geq 5$ , Degradome Data  $\geq 3$ ), we identified five candidate targets for *miR-505-3p* (Figure 4A). To further identify, expression of these targets was estimated in *miR-505-3p*-overexpressed A2780 and OVCAR-3 cells. Compared to other genes, we found that *ZNF326* expression was significantly lowered upon *miR-505-3p* upregulation (Figure 4B). Moreover, RT-qPCR results showed that OC tissues and cells expressed a high level of *ZNF326* (Figure 4C). Furthermore, the complementary base pairs between the *miR-505-3p* seed region and *ZNF326* 3'-UTR were found (Figure 4D). To examine whether *miR-505-3p* could interact with *ZNF326*, we performed a







**Figure 5** KTN1-AS1 regulated OC cell proliferation and invasion by *ZNF326*. (A) *ZNF326* expression in OC cells transfected with pcDNA3.1/*ZNF326* was detected by RT-qPCR. (B) Proliferation of OC cells transfected with indicated plasmids was evaluated by colony formation assay. (C) Flow cytometry analysis was applied to evaluate apoptosis of OC cells with indicated transfection. (D,E) The migration and invasion of OC cells with indicated transfection was determined by wound healing assay and Transwell assay. The percentage of wound healing area and the number of cell clones in different groups at 24 hours were counted. Only use 1% crystal violet staining solution for staining. Photographs were taken using an inverted microscope. The magnification is 100× (wound healing assay) and 400× (transwell assay). \*,  $P < 0.05$ ; \*\*,  $P < 0.01$ ; \*\*\*,  $P < 0.001$ . NC, negative control; OC, ovarian cancer; RT-qPCR, real-time quantitative polymerase chain reaction.

*ZNF326* abolished the inhibited proliferation caused by silencing *KTN1-AS1* (Figure 5B). Results of the flow cytometry analysis revealed that *KTN1-AS1* downregulation could fortify the apoptosis rate of OC cells, but *ZNF326* overexpression impaired the increase in the apoptosis rate (Figure 5C). Furthermore, we discovered that the migration suppressed by *KTN1-AS1* knockdown was restored by transfecting pcDNA3.1/*ZNF326* (Figure 5D). In addition, *ZNF326* overexpression also rescued the inhibitive effect of silenced *KTN1-AS1* on the invasion of OC cells (Figure 5E).

In summary, *KTN1-AS1* increased the *ZNF326* level to enhance OC cell growth.

## Discussion

OC is the main cause of global female cancer incidence and cancer-related mortality (Webb and Jordan, 2017). (1). Most ovarian malignancies originate from the less coelomic epithelium follicular cells and germ cells (26). As the most lethal malignancy in females, only 15% of OC cases are

found at the low-grade stage, and patients with high-grade OC usually relapse within about 16 months (27). Therefore, to prevent or treat OC, we need to find molecular networks that offer prognostic and targeted therapeutic values (28). In recent years, lncRNAs were found to be gene expression modulators of cancer cell phenotypes and are considered promising therapeutic targets for OC treatment (29). Our study confirmed that lncRNA *KTN1-AS1* was upregulated in OC. Functional experiments found that *KTN1-AS1* knockdown slowed the proliferation, migration, and invasion of OC cells, and accelerated apoptosis. This highlighted the oncogenic nature of *KTN1-AS1* in OC.

H3K27ac is known to modify the histone post-translation and transcriptionally activates gene expression by associating with the activity-enhancing regulatory elements (30,31). Existing research has shown that H3K27ac in the promoter region triggered the upregulation of some lncRNAs. For example, H3K27ac in the promoter region of LINC00519 caused the upregulated level of LINC00519 in lung squamous cell carcinoma (32). Furthermore, H3K27ac in the *PLAC2* promoter region activated the expression of *PLAC2* in oral squamous cell carcinoma (33). In this study, we found, through the UCSC website, that the *KTN1-AS1* promoter region was highly enriched with H3K27ac, which was further validated by ChIP assay. The decreased *KTN1-AS1* level by C646 further confirmed that H3K27ac modification caused the upregulation of *KTN1-AS1*. *CBP* is mainly responsible for histone acetylation and gene transcription (25,34). We confirmed that *CBP* bound to the *KTN1-AS1* promoter triggered the H3K27ac, thereby inducing *KTN1-AS1* upregulation.

Regarding mechanisms, lncRNAs were shown as ceRNAs that lncRNAs release mRNAs from posttranscriptional silence induced by miRNAs through sponging miRNAs in cancer progression (35,36). MiRNAs are small noncoding RNAs (18~24 nucleotides) that bind to target mRNA 3'-UTR to suppress protein translation and/or reversely modulate mRNA stability (37). Importantly, miRNAs play significant roles in cancer progression by serving as tumor facilitators or suppressors (38,39). In our research, *miR-505-3p* was identified as a downregulated miRNA and interacted with *KTN1-AS1*. Previously, *miR-505-3p* was shown to be expressed in a low amount and exerted an inhibitive role in pancreatic cancer (40) and breast cancer (41). Moreover, we identified *ZNF326* as the downstream gene of *miR-505-3p*. Former study implied that *ZNF326* increased HDAC7 expression and activated the Wnt pathway to promote a malignant phenotype in glioma (42). In non-small cell lung

cancer, *ZNF326* facilitated cell proliferation by modulating ERCC1 expression (43). Finally, restoration experiments suggested that *KTN1-AS1* enhanced proliferation, migration, and invasion, and attenuated apoptosis by increasing *ZNF326*.

In summary, our study shows that H3K27ac-induced lncRNA *KTN1-AS1* accelerated proliferation, migration, and invasion of OC cells, and impeded apoptosis of OC cells through the *miR-505-3p/ZNF326* axis. These findings provide potential for a new molecular-targeted treatment method for OC patients.

## Acknowledgments

We appreciate each participant for providing assistance in this study.

**Funding:** This work was supported by National Nature Science Foundation of China (No. 82072859) and Guangzhou Science and Technology Program (No. 202102010292) to Kun Shi, Guangzhou Science and Technology Department (No. 202102020059) and Research foundation of Guangzhou Women and Children's Medical Center for Clinical Doctor to Xiaohui Xie.

## Footnote

**Reporting Checklist:** The authors have completed the MDAR reporting checklist. Available at <https://atm.amegroups.com/article/view/10.21037/atm-22-443/rc>

**Data Sharing Statement:** Available at <https://atm.amegroups.com/article/view/10.21037/atm-22-443/dss>

**Conflicts of Interest:** All authors have completed the ICMJE uniform disclosure form (available at <https://atm.amegroups.com/article/view/10.21037/atm-22-443/coif>). The authors have no conflicts of interest to declare.

**Ethical Statement:** The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was approved by the Ethics Committee of Guangzhou Women and Children's Medical Center, Guangzhou Medical University (approval No. 2020-KY-021-03). The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). An informed consent form was signed by each patient.

*Open Access Statement:* This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

## References

1. Webb PM, Jordan SJ. Epidemiology of epithelial ovarian cancer. *Best Pract Res Clin Obstet Gynaecol* 2017;41:3-14.
2. Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. *CA Cancer J Clin* 2017;67:7-30.
3. Matz M, Coleman MP, Carreira H, et al. Worldwide comparison of ovarian cancer survival: Histological group and stage at diagnosis (CONCORD-2). *Gynecol Oncol* 2017;144:396-404.
4. Iyer MK, Niknafs YS, Malik R, et al. The landscape of long noncoding RNAs in the human transcriptome. *Nat Genet* 2015;47:199-208.
5. Batista PJ, Chang HY. Long noncoding RNAs: cellular address codes in development and disease. *Cell* 2013;152:1298-307.
6. Shi X, Sun M, Liu H, et al. Long non-coding RNAs: a new frontier in the study of human diseases. *Cancer Lett* 2013;339:159-66.
7. Liu H, Shang X, Zhu H. LncRNA/DNA binding analysis reveals losses and gains and lineage specificity of genomic imprinting in mammals. *Bioinformatics* 2017;33:1431-6.
8. Furlan G, Rougeulle C. Function and evolution of the long noncoding RNA circuitry orchestrating X-chromosome inactivation in mammals. *Wiley Interdiscip Rev RNA* 2016;7:702-22.
9. Zhang W, Chen L, Wu J, et al. Long noncoding RNA TUG1 inhibits osteogenesis of bone marrow mesenchymal stem cells via Smad5 after irradiation. *Theranostics* 2019;9:2198-208.
10. Kim J, Piao HL, Kim BJ, et al. Long noncoding RNA MALAT1 suppresses breast cancer metastasis. *Nat Genet* 2018;50:1705-15.
11. Zhuo W, Liu Y, Li S, et al. Long Noncoding RNA GMAN, Up-regulated in Gastric Cancer Tissues, Is Associated With Metastasis in Patients and Promotes Translation of Ephrin A1 by Competitively Binding GMAN-AS. *Gastroenterology* 2019;156:676-691.e11.
12. Yan J, Jia Y, Chen H, et al. Long non-coding RNA PXN-AS1 suppresses pancreatic cancer progression by acting as a competing endogenous RNA of miR-3064 to upregulate PIP4K2B expression. *J Exp Clin Cancer Res* 2019;38:390.
13. Hong W, Ying H, Lin F, et al. LncRNA LINC00460 Silencing Represses EMT in Colon Cancer through Downregulation of ANXA2 via Upregulating miR-433-3p. *Mol Ther Nucleic Acids* 2020;19:1209-18.
14. Cao Y, Xiong JB, Zhang GY, et al. Long Noncoding RNA UCA1 Regulates PRL-3 Expression by Sponging MicroRNA-495 to Promote the Progression of Gastric Cancer. *Mol Ther Nucleic Acids* 2020;19:853-64.
15. Wang Q, Yan SP, Chu DX, et al. Silencing of Long Non-coding RNA RP1-93H18.6 Acts as a Tumor Suppressor in Cervical Cancer through the Blockade of the PI3K/Akt Axis. *Mol Ther Nucleic Acids* 2020;19:304-17.
16. Liang H, Yu T, Han Y, et al. LncRNA PTAR promotes EMT and invasion-metastasis in serous ovarian cancer by competitively binding miR-101-3p to regulate ZEB1 expression. *Mol Cancer* 2018;17:119. Erratum in: *Mol Cancer* 2021;20:64.
17. Liu E, Liu Z, Zhou Y. Carboplatin-docetaxel-induced activity against ovarian cancer is dependent on up-regulated lncRNA PVT1. *Int J Clin Exp Pathol* 2015;8:3803-10.
18. Jin Y, Feng SJ, Qiu S, et al. LncRNA MALAT1 promotes proliferation and metastasis in epithelial ovarian cancer via the PI3K-AKT pathway. *Eur Rev Med Pharmacol Sci* 2017;21:3176-84.
19. Gao Y, Meng H, Liu S, et al. LncRNA-HOST2 regulates cell biological behaviors in epithelial ovarian cancer through a mechanism involving microRNA let-7b. *Hum Mol Genet* 2015;24:841-52.
20. Zhang, L. LncRNA KTN1-AS1 promotes tumor growth of hepatocellular carcinoma by targeting miR-23c/ERBB2IP axis. *Biomed Pharmacother* 2019. 109: p. 1140-1147.
21. Li C, Zhao W, Pan X, et al. LncRNA KTN1-AS1 promotes the progression of non-small cell lung cancer via sponging of miR-130a-5p and activation of PDPK1. *Oncogene* 2020;39:6157-71.
22. Hu X, Xiang L, He D, et al. The long noncoding RNA KTN1-AS1 promotes bladder cancer tumorigenesis via KTN1 cis-activation and the consequent initiation of Rho GTPase-mediated signaling. *Clin Sci (Lond)* 2021;135:555-74.
23. Ding G, Li W, Liu J, et al. LncRNA GHET1 activated by

- H3K27 acetylation promotes cell tumorigenesis through regulating ATF1 in hepatocellular carcinoma. *Biomed Pharmacother* 2017;94:326-31.
24. Zhang E, Han L, Yin D, et al. H3K27 acetylation activated-long non-coding RNA CCAT1 affects cell proliferation and migration by regulating SPRY4 and HOXB13 expression in esophageal squamous cell carcinoma. *Nucleic Acids Res* 2017;45:3086-101.
  25. Bose DA, Donahue G, Reinberg D, et al. RNA Binding to CBP Stimulates Histone Acetylation and Transcription. *Cell* 2017;168:135-149.e22.
  26. La Vecchia C. Ovarian cancer: epidemiology and risk factors. *Eur J Cancer Prev* 2017;26:55-62.
  27. Moufarrij S, Dandapani M, Arthofer E, et al. Epigenetic therapy for ovarian cancer: promise and progress. *Clin Epigenetics* 2019;11:7.
  28. Muinao T, Pal M, Deka Boruah HP. Origins based clinical and molecular complexities of epithelial ovarian cancer. *Int J Biol Macromol* 2018;118:1326-45.
  29. Panoutsopoulou K, Avgeris M, Scorilas A. miRNA and long non-coding RNA: molecular function and clinical value in breast and ovarian cancers. *Expert Rev Mol Diagn* 2018;18:963-79.
  30. Wang S, Zang C, Xiao T, et al. Modeling cis-regulation with a compendium of genome-wide histone H3K27ac profiles. *Genome Res* 2016;26:1417-29.
  31. Creighton MP, Cheng AW, Welstead GG, et al. Histone H3K27ac separates active from poised enhancers and predicts developmental state. *Proc Natl Acad Sci U S A* 2010;107:21931-6.
  32. Ye P, Lv X, Aizemaiti R, et al. H3K27ac-activated LINC00519 promotes lung squamous cell carcinoma progression by targeting miR-450b-5p/miR-515-5p/YAP1 axis. *Cell Prolif* 2020;53:e12797.
  33. Chen F, Qi S, Zhang X, et al. lncRNA PLAC2 activated by H3K27 acetylation promotes cell proliferation and invasion via the activation of Wnt/ $\beta$ -catenin pathway in oral squamous cell carcinoma. *Int J Oncol* 2019;54:1183-94.
  34. Dong H, Wang W, Mo S, et al. SP1-induced lncRNA AGAP2-AS1 expression promotes chemoresistance of breast cancer by epigenetic regulation of MyD88. *J Exp Clin Cancer Res* 2018;37:202.
  35. Liu K, Yao H, Wen Y, et al. Functional role of a long non-coding RNA LIFR-AS1/miR-29a/TNFAIP3 axis in colorectal cancer resistance to photodynamic therapy. *Biochim Biophys Acta Mol Basis Dis* 2018;1864:2871-80.
  36. Chen X, Zeng K, Xu M, et al. SP1-induced lncRNA-ZFAS1 contributes to colorectal cancer progression via the miR-150-5p/VEGFA axis. *Cell Death Dis* 2018;9:982.
  37. Ambros V. The functions of animal microRNAs. *Nature* 2004;431:350-5.
  38. Khan MA, Zubair H, Srivastava SK, et al. Insights into the Role of microRNAs in Pancreatic Cancer Pathogenesis: Potential for Diagnosis, Prognosis, and Therapy. *Adv Exp Med Biol* 2015;889:71-87.
  39. Dong P, Xiong Y, Yue J, et al. miR-34a, miR-424 and miR-513 inhibit MMSET expression to repress endometrial cancer cell invasion and sphere formation. *Oncotarget* 2018;9:23253-63.
  40. Xu Z, Zhang D, Zhang Z, et al. MicroRNA-505, Suppressed by Oncogenic Long Non-coding RNA LINC01448, Acts as a Novel Suppressor of Glycolysis and Tumor Progression Through Inhibiting HK2 Expression in Pancreatic Cancer. *Front Cell Dev Biol* 2021;8:625056.
  41. Zhao P, Guan H, Dai Z, et al. Long noncoding RNA DLX6-AS1 promotes breast cancer progression via miR-505-3p/RUNX2 axis. *Eur J Pharmacol* 2019;865:172778.
  42. Yu X, Wang M, Wu J, et al. ZNF326 promotes malignant phenotype of glioma by up-regulating HDAC7 expression and activating Wnt pathway. *J Exp Clin Cancer Res* 2019;38:40. Erratum in: *J Exp Clin Cancer Res* 2020;39:14.
  43. Wu J, Zhang X, Han Q, et al. ZNF326 promotes proliferation of non-small cell lung cancer cells by regulating ERCC1 expression. *Lab Invest* 2019;99:169-79.

(English Language Editor: C. Mullens)

**Cite this article as:** Xie X, Wen Q, Yang X, Chen W, Liu Y, Liu W, Zhang T, Xu C, Shi K. H3K27ac-activated lncRNA KTN1-AS1 aggravates tumor progression by miR-505-3p/ZNF326 axis in ovarian cancer. *Ann Transl Med* 2022;10(10):599. doi: 10.21037/atm-22-443