

Michigan Medicine Tracheostomy Guidelines in COVID-19 Era:

Based on our current understanding of COVID19, it is spread mainly through from person-to-person either via close contact with one another (within about 6 feet) or through respiratory droplets produced when an infected person coughs or sneezes. Procedures such as tracheostomy and laryngectomy can result in significant aerosol generation and lead to increased risk to others.

In cases of airway intervention, the goal is to try minimize aerosolization and risk of viral transmission to health care providers.

Performance of Tracheostomy is considered a **high-risk aerosol generating procedure (AGP)**. **Avoid performing tracheostomy in COVID19/PUI patients if possible** to avoid increased aerosolization risks to health care providers.

In an attempt to minimize the exposure to COVID-19 to healthcare workers, the following measures are suggested in tracheostomy (or laryngectomy) patients with either a confirmed diagnosis of COVID-19, persons under investigation (PUI), or unknown/asymptomatic status.

We will continue to follow all guidelines from the Michigan Medicine and Infection Prevention and Epidemiology (IPE). This is a dynamic time and best practices may change, so please defer to updated official communications from Michigan Medicine.

Emergency Trach/Airway:

Indication: Emergency with impending loss of airway where transoral ETT placement is unsuccessful and cannot be safely performed. If stable, attempt to transfer to the OR.

At this time, in patients with unknown COVID19 status, COVID19-positive or PUI, all health care providers performing emergent tracheostomy should wear COVID19 PPE.

In the setting of an emergency airway, the surgical airway team should don appropriate COVID19 PPE (see Michigan Medicine guidelines for the use of PPE) **before entering the room** and assisting with the airway and patient care.

Personal Protective Equipment for COVID19

Please note that institutional guidelines for PPE use are subject to change. Updated guidance can be accessed at-

<http://www.med.umich.edu/i/covid19-careplan/protect.html>

- PPE should be at the door/unit of all COVID19 and PUI patients
- **PPE should be donned appropriately PRIOR TO ENTERING THE ROOM before proceeding to assist**

Appropriate PPE is critical based on the Viral Exposure Risk:

- The Covid-19 virus is found in the respiratory tract and secretions, the GI tract and secretions, as well as blood. Eye protection is critically important.
- Though data are sparse, the virus may potentially be aerosolized during the use of electrocautery and during airway manipulation. **Hence PAPR/N-95 use for all tracheostomy related cases and procedures.** (<http://www.med.umich.edu/i/covid19-careplan/protect.html>)

Unprotected healthcare personnel SHOULD NOT be allowed in a room where an aerosol-generating procedure is being or has been conducted.

If COVID19 status is unknown, manage patient as if they are COVID-19 positive. Given respiratory symptoms they will fulfil criteria for suspected COVID-19 and there will not be time for testing in this type of emergency situation.

End of procedure:

• **Careful DOFFING is critically important.** This is the moment of highest risk for self-contamination. Proper removal of PPE is critical to avoid cross contamination & potential exposure

<http://www.med.umich.edu/i/covid19-careplan/protect.html>

In known COVID or PUI patients, the room will need to be cleared according to institutional guidelines (see link below) to allow adequate air clearance.

http://www.med.umich.edu/i/ice/resources/coronavirus/air_clearance_by_room.pdf

Elective Tracheostomy in non-ICU setting:

Elective tracheostomy may be necessary in patients who are not critically ill, (ie: major head and neck cancer/reconstructive cases to bypass potential airway obstruction, non-emergent subglottic/tracheal stenosis, etc)

If possible, avoid tracheostomy or delay the procedure as long as avoidance or delay would not cause undue harm to the patient or require use of other scarce resources (ie ICU bed). Decisions to perform pre-operative COVID testing should be based on institutional guidelines.

Protocol for pre-op testing and algorithm for determination of need to proceed to OR:

http://www.med.umich.edu/i/ice/resources/coronavirus/ppe_or_guidance.pdf

Elective Tracheostomy for Mechanically Ventilated Patients in the ICU:

Carefully consider timing and indication of tracheostomy in all patients who being considered for tracheostomy, particularly those that are COVID19 positive or PUI. Given the high risk of aerosol generation both during tracheostomy and after the procedure, defer tracheostomy in the COVID19/PUI population until absolutely necessary.

In COVID-19 positive patients and considering elective tracheostomy:

1. Confirm improvement of symptoms and stable pulmonary status
2. Wait at least 3 weeks from intubation (allows for optimal viral clearance)
Extenuating circumstances may present where tracheostomy may be required earlier than 3 weeks; in these exceptional situations, the clinical benefits of tracheostomy must be carefully weighed against the high transmission risk in COVID-19 tracheostomy patients and should be done in a multidisciplinary fashion.

Decision for tracheostomy in any ICU patient should be a multi-disciplinary decision as transmission risk in a COVID-19 patient with a tracheostomy is high.

PPE for Elective Tracheostomy: PPE appropriate for an AGP should be used, per institutional guidelines.

Type and Location of Tracheostomy Procedures:

- There is no high quality data on the degree of aerosolization with open versus percutaneous tracheostomy techniques
- The use of bronchoscopy with percutaneous tracheostomy may increase the risk of aerosolization during the procedure, but the degree of aerosolization and risk to the health care worker is felt to be equivalent to open procedures. *Therefore, can consider either open tracheostomy or percutaneous tracheostomy at the discretion of the faculty performing the procedure.*

- For percutaneous tracheostomy performed with alternate techniques, such as with real-time ultrasound guidance without bronchoscopy, will further minimize risk to providers
- *Location of tracheostomy:* Perform tracheostomy (either percutaneous or open) in the ICU if possible.
 - Tracheostomy is favored to be performed in the ICU
 - **Perform tracheostomy in a negative pressure room**; secondary choice is an externally vented room with HEPA filter if airborne room is unavailable. Avoid performing tracheostomy in neutral pressure rooms
 - In cases not felt to be safe for bedside tracheostomy in the ICU, then transport to the OR is appropriate.
 - Follow Institutional protocols for transport of COVID19/PUI patients (http://www.med.umich.edu/i/ice/resources/coronavirus/patient_transport.pdf)
- If a known COVID or PUI patient is taken to the OR, the room will need to be cleared for 20-35 minutes to allow adequate air clearance. (http://www.med.umich.edu/i/ice/resources/coronavirus/air_clearance_by_room.pdf)

In previously asymptomatic patients who are deemed COVID-19-NEGATIVE based on testing within 48 hours of the planned procedure, standard PPE is appropriate.

Unique Steps/Precautions During Tracheostomy Procedure:

These are unique and critical steps within a standard tracheostomy for COVID19/PUI patients:

- Preparation:
 - The most senior/experienced surgeon/proceduralist available should perform the tracheostomy in COVID19/PUI patients
 - Limit personnel in the room to individuals essential for safe completion of the procedure
 - Maintain neuromuscular blockade throughout the procedure
 - Consider IV Glycopyrrolate 0.4mg to reduce secretions
 - Inject with 2% lidocaine with 1:100,000 epinephrine to minimize bleeding and need for suctioning
 - Runner outside room
 - Ensure HEPA viral filter on ventilator and suctioning
 - Double gloving
 - Full draping to minimize contamination of surfaces
 - Impervious sheets around patients
 - Avoid electrocautery if possible; strongly recommend smoke evacuator cautery
 - If possible, use single-use bronchoscope for percutaneous trach
- Use the following steps to minimize aerosolization in open tracheostomy or percutaneous:
 - During open tracheostomy:
 - Anesthesia informed of imminent tracheal incision; all team members are prepared
 - Preoxygenation 100% for 3min then apnea
 - Ventilator 'OFF'
 - Cuff deflated just before incision down to trachea or pushed distal to avoid accidentally popping balloon of the endotracheal tube (ETT)
 - ETT Pulled back 3 cm and visualization of tip of ETT at tracheotomy
 - **Minimize tracheal suctioning to avoid aerosolization**
 - Resume mechanical ventilation only **after** the tracheostomy tube balloon is inflated and a closed circuit re-established; confirm HME filter is connected to tracheostomy tube
 - After removal of ETT, place ETT in plastic bag for disposal

- During percutaneous tracheostomy:
 - To minimize aerosolization during withdrawal of the endotracheal tube:
 - Confirm that patient has been on FiO2 100% at least 5 minutes prior to the tracheostomy and continues throughout the procedure
 - The proceduralist must notify other members of the team that withdrawal of the tube will commence
 - Turn off mechanical ventilation
 - Withdraw endotracheal tube under appropriate guidance, consider Doppler to determine airflow as ETT pulled back
 - After extubation, place ETT in plastic bag for disposal
 - Inflate cuff, then resume mechanical ventilation
 - To minimize aerosolization during dilation:
 - Confirm that patient has been on FiO2 100% through the procedure
 - The proceduralist must notify other members of the team that dilatation will commence
 - Turn off mechanical ventilation
 - Perform dilatation and delivery of tracheostomy tube, cover stoma with gauze between these steps
 - Consider performing dilatation and tube delivery under a clear plastic drape
 - Resume mechanical ventilation only **after** the tracheostomy tube balloon is inflated and a closed circuit re-established
- End of procedure:
 - Consider a **xeroform trach pants** to further seal the tracheostomy incision to prevent leak and aerosolization until the stoma has closed around the tracheostomy tube
 - Need to ensure adequate cuff-pressure immediately to minimize leaks
 - Use the largest tracheostomy tube appropriate for the patient to allow bronchoscopy in ICU patients; do not use fenestrated tracheostomy tube

Care of Tracheostomy Patients:

PPE for care of tracheostomy: http://www.med.umich.edu/i/ice/resources/coronavirus/ppe_pt_care.pdf

COVID19/PUI/Symptomatic Patients:

Because tracheostomy and tracheal suctioning is considered an AGP, all COVID19/PUI patients not connected to mechanical ventilation **should have an HME** (heat and moisture exchanger)/filter fitted to the tracheostomy tube.

- **Closed suctioning if available is strongly recommended**
- During suctioning, change of inner cannula, disconnect from ventilator circuit, bronchoscopy through tracheostomy tube, and tracheostomy tube change, the caregiver(s) must wear appropriate COVID19 PPE (as above)
- Keeping tracheostomy tube inflated to minimize secretions from around the tube
- **Avoid routine tracheostomy tube exchange** unless absolutely necessary or planning for decanulation. Tracheostomy tube exchange is considered a high risk AGP.

Laryngectomy patients are a unique group of patients that also have similar AGP risks to health care workers with generation of aerosolized respiratory droplets. Laryngectomy HME should be placed immediately and kept at all times regardless of COVID19 status given many carriers are asymptomatic. In COVID19/PUI laryngectomy patients, the addition of appropriate PPE based on their status should be in place for all suctioning, laryngectomy tube exchange, or airway manipulation.

http://www.med.umich.edu/i/ice/resources/coronavirus/ppe_pt_care.pdf

Untested/Asymptomatic Patients:

In untested/asymptomatic tracheostomy and laryngectomy patients should have an HME/filter on at all times and if possible/available, closed suctioning is recommended. *Strongly consider testing these patients if symptoms develop and placement in special pathogen precautions while awaiting testing results*; however, unless testing COVID19-positive, follow standard precautions (http://www.med.umich.edu/i/ice/resources/coronavirus/ppe_pt_care.pdf) with any tracheostomy/stomal interventions (i.e.: suctioning, inner cannula exchange, changing HME/filter).

- **Avoid routine tracheostomy tube exchange** unless absolutely necessary or planning for decannulation
- **Tracheostomy tube exchange is considered a high risk AGP**, thus COVID-19 PPE is recommended (even in asymptomatic/untested patients), limit the number of providers to only those essential for the procedure, and must communicate/coordinate with nursing as the room will need to be cleared for 60 minutes after the tracheostomy tube exchange (http://www.med.umich.edu/i/ice/resources/coronavirus/ppe_pt_care.pdf)

Decannulation: Should be done as soon as medically safe and immediately place occlusive dressing. Providers should be aware this can take up to 2 weeks to close and in 1-3% of patients will not close.

Laryngectomy patients cannot be decannulated nor should occlusive dressing be placed.

Special Category: Care of Untested/Asymptomatic Pediatric Tracheostomy Patients:

- Pediatric tracheostomy patients who are asymptomatic should be maintained in standard precautions with healthcare providers wearing the appropriate PPE (http://www.med.umich.edu/i/ice/resources/coronavirus/ppe_pt_care.pdf).
- Patients should use closed suctioning systems for trach tube suctioning whenever possible to limit circuit disconnection and decrease risk of exposure
- Patients should use HME during the day and if possible at night/naps as much as tolerated
 - If HME is not tolerated, can use of trach mask or tracheostomy covers with the goal to cover the tracheostomy as much as possible to minimize droplet spread
- In pediatric patients, routine tracheostomy tube exchanges should occur every 4 weeks unless there is concern for mucus plugging or emergency situation. *Tracheostomy tube exchange should absolutely be performed if there is concern for airway patency.*
- Tracheostomy tube exchanges performed by health care providers should be done with PPE according to institutional guidelines (http://www.med.umich.edu/i/ice/resources/coronavirus/ppe_pt_care.pdf).
- Families can continue to participate in tracheostomy tube care in the hospital as appropriate. Tracheostomy tube exchanges with family caregivers will be coordinated by Peds OTO Trach Team. Per IPE, family caregivers should wear loop mask only when participating in tracheostomy tube exchanges.

Acknowledgements: MM Tracheostomy Working group (Steven B. Chinn, MD MPH; Jose L. De Cardenas, MD; Sean P. Edwards, DDS MD; Jennifer Helman, NP; Kelly Krawke, RN; William R. Lynch, MD; Kelly M. Malloy, MD; Pauline K. Park, MD; Venkatakrishna Rajajee, MD; Samuel A. Schechtman, MD; Marc C. Thorne, MD; Laraine L. Washer, MD)

References:

1. Wei WI, et al. Safe Tracheostomy for patients with Severe Acute Respiratory Syndrome. Laryngoscope, 2003; PMID: 14520105
2. Tay JK, Khoo ML, Loh WS. Surgical Considerations for Tracheostomy During the COVID-19 Pandemic: Lessons Learned From the Severe Acute Respiratory Syndrome Outbreak. *JAMA Otolaryngol Head Neck Surg*. Published online March 31, 2020. doi:10.1001/jamaoto.2020.0764
3. Vargas M, Servillo, G. Improving staff safety during tracheostomy in COVID-19 patients. *Head and Neck*. Published online April 9, 2020.

4. Heyd, CP. Tracheostomy protocols during COVID-19 outbreak. COVID vs Cancer: Impact on Head and Neck Oncology. Head and Neck. Preprint April 8, 2020.
5. UKENT Guidelines: <https://www.entuk.org/tracheostomy-guidance-during-covid-19-pandemic>
6. AAO-HNS Guidelines: <https://www.entnet.org/content/tracheotomy-recommendations-during-covid-19-pandemic>
7. Brewster DJ, et al. Consensus Statement: Safe airway society principles of airway management and tracheal intubation specific to COVID-19 adult patient group. Medical Journal OF Australia. 2020.
8. Zhonghua et al. Expert consensus on preventing nosocomial transmission during respiratory care for critically ill patients infected by 2019 novel coronavirus pneumonia. Respiratory care committee of Chinese Thoracic Society. 2020. PMID: 32077661.
9. icmanaesthesiacovi-19.org
10. ENT UK Guidance for ENT during the COVID-19 pandemic
11. van Doremalen N, Bushmaker T, Morris DH et al. Aerosol and surface stability of HCoV-19 (SARS-CoV-2) compared to SARS-CoV-1. NEJM in press doi: <https://doi.org/10.1101/2020.03.09.20033217>.
The <https://www.medrxiv.org/content/10.1101/2020.03.09.20033217v1.full.pdf>
12. Tran K, Cimon K, Severn M, Pessoa-Silva CL, Conly J. Aerosol generating procedures and risk of transmission of acute respiratory infections to healthcare workers: a systematic review. PLoS One 2012; 7: e35797
13. Public Health England. Environmental decontamination, in COVID-19: infection prevention and control guidance (<https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control/wuhan-novel-coronavirus-wn-cov-infection-prevention-and-control-guidance#decon>)
14. Loeb M, McGeer A, Henry B, et al. SARS among critical care nurses, Toronto. Emerg Infect Dis 2004; 10: 251-5
15. COVID-19 Guidance for Infection Prevention and Control in Healthcare Settings. Version 1.0.
16. ICU Management and Practice. China with the World: COVID 19 Experts Dialogues (<https://healthmanagement.org/c/icu/pressrelease/china-with-the-world-covid-19-experts-dialogues-the-2nd-talk-transcript>)
17. Liew MF et al. Preparing for COVID-19: early experience from an intensive care unit in Singapore. Crit Care. 2020 PMID: 32151274
18. Chan JYK et al. Practical Aspects of Otolaryngologic Clinical Services During the 2019 Novel Coronavirus Epidemic: An Experience in Hong Kong. 2020. doi:10.1001/jamaoto.2020.0488.
19. Wölfel R, Corman, V.M., Guggemos, W. et al. Virological assessment of hospitalized patients with COVID-2019. *Nature* (2020). <https://doi-org.proxy.lib.umich.edu/10.1038/s41586-020-2196-x>
20. Yang X, et al.. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. Lancet Respir Med. 2020 Feb 24. pii: S2213-2600(20)30079-5. doi: 10.1016/S2213-2600(20)30079-5. [Epub ahead of print]
21. Zou L, Ruan F, Huang M, et al. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. N Engl J Med 2020; published online Feb 19. DOI:10.1056/NEJMc200173.
22. Lescure F-X, Bouadma L, Nguyen D, et al. Clinical and virological data of the first cases of COVID-19 in Europe: a case series. Lancet Infect Dis 2020; published online March 27. [https://doi.org/10.1016/S1473-3099\(20\)30200-0](https://doi.org/10.1016/S1473-3099(20)30200-0).